

IB/2004/01723



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

REC'D 16 JUN 2004

WIPO

PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

03405365.2 ✓

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk

BEST AVAILABLE COPY



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

Anmeldung Nr.:
Application no.: 03405365.2 ✓
Demande no:

Anmeldetag:
Date of filing: 23.05.03 ✓
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Aponetics AG
Benkenstrasse 254
4108 Witterswil
SUISSE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Furazanobenzimidazoles

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

C07D271/00

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PT RO SE SI SK TR LI

Furazanobenzimidazoles

The invention relates to novel substituted furazanobenzimidazoles, processes for the preparation thereof, pharmaceutical compositions containing same, the use thereof
5 optionally in combination with one or more other pharmaceutically active compounds for the therapy of neoplastic diseases and autoimmune diseases, and a method for the treatment of such a diseases.

Background of the invention

10

Cancer is one of the leading causes of death in humans. Although a variety of drugs against neoplastic diseases have been developed and techniques are available such as surgery and radiation therapy, there is still a need for alternative and improved methods of
treatment of neoplastic diseases.

15

Autoimmune diseases are associated with abnormal lymphoproliferation as a result of defects in the termination of lymphocyte activation and growth. Often, such diseases are associated with inflammation like rheumatoid arthritis, insulin dependent diabetes mellitus, multiple sclerosis, systemic lupus erythematosus and the like. The treatment of such
20 diseases is focused on anti-inflammatory and immunosuppressive drugs which in numerous cases show severe side effects. Hence, there is a need for alternative drugs with a new mode of action showing less side effects.

25

Apoptosis is a term used to describe a series of cellular events which occur to bring about programmed cell death. There are various apoptotic pathways, some of which have been characterized, whereas others remain to be elucidated. If the balance between cell division and apoptosis is disturbed, life-threatening diseases including cancer, autoimmune disorders, neurodegenerative and cardiovascular diseases may occur.

30

In recent years it has become evident that programmed cell death (apoptosis) is as important to the health of a multicellular organism as cell division. By repeated cell division and differentiation throughout development or tissue repair, surplus or even harmful cells are generated. In order to maintain tissue homeostasis these cells have to be removed or killed. The delicate interplay between cell growth and apoptosis in an organism is mirrored

in the complex molecular balance that determines whether an individual cell undergoes division, arrests in the cell-cycle-or-commits to programmed cell death.

5 Dysregulation of cell proliferation, or lack of appropriate cell death, has wide ranging clinical implications. A number of diseases associated with such dysregulation involve hyperproliferation, inflammation, tissue remodeling and repair. Familiar indications in this category include cancers, restenosis, neointimal hyperplasia, angiogenesis, endometriosis, lymphoproliferative disorders, transplantation related pathologies (graft rejection), polyposis, loss of neural function in the case of tissue remodeling and the like.
10 Such cells may lose the normal regulatory control of cell division, and may also fail to undergo appropriate cell death.

As apoptosis is inhibited or delayed in most types of proliferative, neoplastic diseases, induction of apoptosis is an option for treatment of cancer, especially in cancer types
15 which show resistance to classic chemotherapy, radiation and immunotherapy (Apoptosis and Cancer Chemotherapy, Hickman and Dive, eds., Blackwell Publishing, 1999). Also in autoimmune and transplantation related diseases and pathologies compounds inducing apoptosis may be used to restore normal cell death processes and therefore can eradicate the symptoms and might cure the diseases. Further applications of compounds
20 inducing apoptosis may be in restenosis, i.e. accumulation of vascular smooth muscle cells in the walls of arteries, and in persistent infections caused by a failure to eradicate bacteria- and virus-infected cells. Furthermore, apoptosis can be induced or re-established in epithelial cells, in endothelial cells, in muscle cells, and in others which have lost contact with extracellular matrix. These cells are potentially able to colonize
25 other organs and therefore can develop into pathologies like neoplasias, endometriosis and the like.


Summary of the invention

30 Furazanobenzimidazoles of formula (I) are selectively inducing apoptosis in cancer cells, and can be used for the treatment of neoplastic and autoimmune diseases. The invention relates to compounds of formula (I) for use as medicaments as defined hereinafter, to novel compounds of formula (I), to methods of synthesis of such compounds, to pharmaceutical compositions containing compounds of formula (I), to the use of a
35 compounds of formula (I) for the preparation of a pharmaceutical composition for the

treatment of neoplastic and autoimmune diseases, and to methods of treatment of neoplastic and autoimmune diseases using such compounds of formula (I) or of pharmaceutical compositions containing same.

5 Detailed description of the invention

The invention relates to compounds of formula (I)



10

wherein

R represents aryl or heteroaryl optionally substituted by up to four substituents independently selected from

15 alkyl, cycloalkyl, cycloalkyl-lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, lower alkoxy-lower alkoxy-lower alkyl, halo-lower alkoxy-lower alkyl, acyloxy-lower alkyl, heterocyclyl, heterocyclyl-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl, optionally substituted alkenyl, optionally substituted alkynyl, hydroxy, lower alkoxy, optionally substituted alkenyloxy, optionally substituted alkynyloxy, 20 cycloalkoxy, halo-lower alkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, heterocycliloxy, heterocyclyl-lower alkoxy, optionally substituted phenyloxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy, optionally substituted heteroaryl-lower alkoxy, sulfamoyloxy, carbamoyloxy, lower alkylcarbonyloxy,

25 amino, monoalkylamino, dialkylamino, aminocarbonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylcarbonylamino wherein heterocyclyl is bound via a nitrogen atom, aminosulfonylamino wherein each of the two amino groups is optionally substituted by

alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylsulfonylamino wherein heterocyclyl is bound via a nitrogen atom, lower alkoxycarbonylamino, lower alkylcarbonylamino wherein alkyl is optionally substituted by one or two substituents selected from optionally substituted phenyl, guanidyl, halogen, cyano, alkoxy, optionally substituted phenoxy, alkylmercapto and optionally substituted amino; lower alkenylcarbonylamino wherein alkenyl is optionally substituted by one or two substituents selected from lower alkyl, halo-lower alkyl, optionally substituted phenyl, halogen, cyano, alkoxy and optionally substituted amino; amino-lower alkyl or amino-lower alkylamino, wherein the nitrogen atom is unsubstituted or substituted by one or two substituents selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl, lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclylcarbonyl, carboxy, lower alkoxycarbonyl, hydroxy-lower alkoxycarbonyl, lower alkoxy-lower alkoxycarbonyl, optionally substituted phenyl-lower alkoxycarbonyl, cyano, lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl, halo-lower alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, and nitro; and wherein two adjacent substituents together with the atoms of aryl or heteroaryl may form a 5 or 6 membered carbocyclic or heterocyclic ring;

X represents a bond; oxygen; a group C=Y, wherein Y stands for oxygen, nitrogen substituted by hydroxy, alkoxy or optionally substituted amino; an ethylenediyl group optionally substituted by alkyl; or a group CR⁷R⁸;

R¹ and R², independently of each other, represent hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, optionally substituted arylalkyl, optionally substituted heteroarylalkyl, hydroxyalkyl, alkoxyalkyl, hydroxyalkoxyalkyl, alkoxyalkoxyalkyl, cyanoalkyl, optionally substituted alkenyl, optionally substituted alkynyl, or lower alkylcarbonyl wherein lower alkyl is optionally substituted by one or two substituents selected from aryl, optionally substituted amino, alkoxy and aryloxy,

R^3 , R^4 , R^5 and R^6 , independently of each other, represent hydrogen, lower alkyl, halo-lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower-alkoxy-lower alkyl, lower alkoxy-lower alkoxy-lower alkyl, halo-lower alkoxy-lower alkyl, heterocyclyl, heterocyclyl-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl, optionally substituted alkenyl, optionally substituted alkynyl, hydroxy, lower alkoxy, halo-lower alkoxy, cycloalkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, heterocyclyloxy, heterocyclyl-lower alkoxy, optionally substituted phenyloxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy, optionally substituted heteroaryl-lower alkoxy, amino, carbamoyl, sulfamoyl, amino-lower alkyl or amino-lower alkylamino, wherein in each case the nitrogen atom is unsubstituted or substituted by one or two substituents selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl, lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclylcarbonyl, carboxy, lower alkoxycarbonyl, hydroxy-lower alkoxycarbonyl, lower alkoxy-lower alkoxycarbonyl, optionally substituted phenyl-lower alkoxycarbonyl, cyano, lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl, halo-lower alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, or nitro, or R^3 and R^4 , R^4 and R^5 , or R^5 and R^6 together with the atoms of the phenyl ring form a 5 or 6 membered carbocyclic or heterocyclic ring;

R^7 represents hydrogen, lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, lower alkenyl, lower alkynyl, optionally substituted phenyl, lower alkoxy, lower alkenyloxy, lower alkynyloxy; R^8 represents hydrogen, lower alkyl, hydroxy, lower alkoxy or lower alkenyloxy, or R^7 and R^8 together with the carbon they are bound to form a 5 or 6 membered carbocyclic or heterocyclic ring;

and salts thereof; for use as medicaments.

The general terms used hereinbefore and hereinafter preferably have within the context of this disclosure the following meanings, unless otherwise indicated:

5 The prefix "lower" denotes a radical having up to and including a maximum of 7, especially up to and including a maximum of 4 carbon atoms, the radicals in question being either linear or branched with single or multiple branching.

10 Where the plural form is used for compounds, salts, and the like, this is taken to mean also a single compound, salt, or the like.

Double bonds in principle can have E- or Z-configuration. The compounds of this invention may therefore exist as isomeric mixtures or single isomers. If not specified both isomeric forms are intended.

15 Any asymmetric carbon atoms may be present in the (R)-, (S)- or (R,S)-configuration, preferably in the (R)- or (S)-configuration. The compounds may thus be present as mixtures of isomers or as pure isomers, preferably as enantiomer-pure diastereomers.

20 The invention relates also to possible tautomers of the compounds of formula (I).

Alkyl has from 1 to 12, preferably from 1 to 7 carbon atoms, and is linear or branched. Alkyl is preferably lower alkyl.

25 Lower alkyl has 1 to 4 carbon atoms and is butyl, such as n-butyl, sec-butyl, isobutyl, tert-butyl, propyl, such as n-propyl or isopropyl, ethyl or methyl. Preferably lower alkyl is methyl or ethyl.

30 Cycloalkyl has preferably 3 to 7 ring carbon atoms, and may be unsubstituted or substituted, e.g. by lower alkyl or lower alkoxy. Cycloalkyl is, for example, cyclohexyl, cyclopentyl, or methylcyclopentyl.

35 Aryl stands for a mono- or bicyclic fused ring aromatic group with 5 to 10 carbon atoms, such as phenyl, 1-naphthyl or 2-naphthyl, or also a partially saturated bicyclic fused ring comprising a phenyl group, such as indanyl, dihydro- or tetrahydronaphthyl.

In optionally substituted phenyl, substituents are preferably lower alkyl, lower alkoxy, lower alkoxy-lower alkoxy, methylenedioxy, halo-lower alkyl, lower alkoxy-lower-alkyl, halo, or nitro.

- 5 Heteroaryl represents an aromatic group containing at least one heteroatom selected from nitrogen, oxygen and sulfur, and is mono- or bicyclic. Monocyclic heteroaryl includes 5 or 6 membered heteroaryl groups containing 1, 2, 3 or 4 heteroatoms selected from nitrogen, sulfur and oxygen. Bicyclic heteroaryl includes 9 or 10 membered fused-ring heteroaryl groups. Examples of heteroaryl include pyrrolyl, thienyl, furyl, pyrazolyl, imidazolyl, triazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiazolyl, isothiazolyl, thiadiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, benzo fused derivatives of such monocyclic heteroaryl groups, such as indolyl, benzimidazolyl or benzofuryl, quinolinyl, isoquinolinyl, quinazolinyl, or purinyl.

- 15 In optionally substituted heteroaryl, substituents are preferably lower alkyl, lower alkoxy, lower alkoxy-lower alkoxy, amino, optionally substituted by one or two substituents selected from lower alkyl, lower alkenyl and alkylcarbonyl, halo-lower alkyl, lower alkoxy-lower alkyl, halo, or nitro.

- 20 Alkenyl contains one or more, e.g. two or three, double bonds, and is preferably lower alkenyl, such as 1- or 2-butenyl, 1-propenyl, allyl or vinyl.

Alkynyl is preferably lower alkynyl, such as propargyl or acetylenyl.

- 25 Ethylenediyl designates a vinyl group bound to R and to methylene as defined in formula (I). The bonds to R and to methylene may be in geminal or vicinal position of the vinyl group.

- 30 In optionally substituted alkenyl or alkynyl, substituents are preferably lower alkyl, lower alkoxy, halo or di(lower alkyl)amino, and are connected with a saturated carbon atom of alkenyl or alkynyl or with an unsaturated carbon atom of alkenyl.

- 35 Heterocyclyl designates preferably a saturated, partially saturated or unsaturated, mono- or bicyclic ring containing 4-10 atoms comprising one, two or three heteroatoms selected from nitrogen, oxygen and sulfur, which may, unless otherwise specified, be carbon or

nitrogen linked, wherein a ring nitrogen atom may optionally be substituted by a group selected from lower alkyl, amino-lower-alkyl, aryl, aryl-lower-alkyl and acyl, and a ring carbon atom may be substituted by lower alkyl, amino-lower alkyl, aryl, aryl-lower alkyl, heteroaryl, lower alkoxy, hydroxy or oxo. Examples of heterocyclyl are pyrrolidinyl, oxazolidinyl, thiazolidinyl, piperidinyl, morpholinyl, piperazinyl, dioxolanyl and tetrahydropyranyl.

Acyl designates, for example, alkylcarbonyl, cyclohexylcarbonyl, arylcarbonyl, aryl-lower alkylcarbonyl, or heteroarylcarbonyl. Lower acyl is preferably lower alkylcarbonyl, in particular propionyl or acetyl.

Hydroxyalkyl is especially hydroxy-lower alkyl, preferably hydroxymethyl, 2-hydroxyethyl or 2-hydroxy-2-propyl.

Cyanoalkyl designates preferably cyanomethyl and cyanoethyl.

Haloalkyl is preferably fluoroalkyl, especially trifluoromethyl, 3,3,3-trifluoroethyl or pentafluoroethyl.

Halogen is fluorine, chlorine, bromine, or iodine.

Lower alkoxy is especially methoxy, ethoxy, isopropoxy, or tert-butoxy.

Arylalkyl includes aryl and alkyl as defined hereinbefore, and is e.g. benzyl, 1-phenethyl or 2-phenethyl.

Heteroarylalkyl includes heteroaryl and alkyl as defined hereinbefore, and is e.g. 2-, 3- or 4-pyridylmethyl, 1- or 2-pyrrolylmethyl, 1-pyrazolylmethyl, 1-imidazolylmethyl, 2-(1-imidazolyl)ethyl or 3-(1-imidazolyl)propyl.

Two adjacent substituents which together with the atoms of aryl or heteroaryl may form a 5 or 6 membered carbocyclic or heterocyclic ring are, for example, propylene, 1- or 2-oxopropylene, 1- or 2-oxapropylene, 1-oxapropylidene, methylenedioxy, difluoromethylenedioxy, 1- or 2-azapropylene, 1- or 2-azapropylidene, 1,2- or 1,3-diazapropylidene, 1,3-diaza-2-oxopropylene, butylene, 1- or 2-oxabutylene, ethylenedioxy,

1- or 2-azabutylene, or 1- or 2-azabutadienyldiene, or such groups carrying further substituents as defined hereinbefore.

5 A 5 or 6 membered carbocyclic or heterocyclic ring formed by substituents R^7 and R^8 together with the carbon atom they are bound to is e.g. cyclopentane, cyclohexane, such rings wherein one or preferably two carbon atoms are replaced by oxygen, or such rings wherein one carbon atom is replaced by oxygen and another one by nitrogen, and is optionally further substituted by lower alkyl, lower alkoxy or lower alkoxy-lower alkyl. Preferred examples are cyclic acetals formed from a carbonyl group with ethylene glycol or monoalkylated glycerin, i.e. rings wherein the substituents R^7 and R^8 together represent 10 1,2-ethylenedioxy or 3-alkoxypropylene-1,2-dioxy.

In substituted amino, the substituents are preferably those mentioned as substituents R^5 and R^6 . In particular, substituted amino is alkylamino, dialkylamino, optionally substituted 15 arylamino, optionally substituted arylalkylamino, lower alkylcarbonylamino, lower alkoxy carbonylamino or optionally substituted aminocarbonylamino.

When X represents a group $C=Y$, wherein Y stands for nitrogen substituted by hydroxy, this corresponds to an oxime function. Oximes and the corresponding oxime alkyl ethers 20 (nitrogen substituted by alkoxy) may be present in E or Z form, or as mixture of isomers. In groups wherein Y stand for nitrogen substituted by optionally substituted amino, this group corresponds to an optionally substituted hydrazone function. Substituents are those considered for substituted amino above, in particular alkylamino, dialkylamino, optionally substituted arylamino or optionally substituted aralkylamino.

25 Salts are especially the pharmaceutically acceptable salts of compounds of formula (I).

Such salts are formed, for example, as acid addition salts, preferably with organic or inorganic acids, from compounds of formula (I) with a basic nitrogen atom, especially the 30 pharmaceutically acceptable salts. Suitable inorganic acids are, for example, halogen acids, such as hydrochloric acid, sulfuric acid, or phosphoric acid. Suitable organic acids are, for example, carboxylic, phosphonic, sulfonic or sulfamic acids, for example acetic acid, propionic acid, octanoic acid, decanoic acid, dodecanoic acid, glycolic acid, lactic acid, fumaric acid, succinic acid, adipic acid, pimelic acid, suberic acid, azelaic acid, malic acid, tartaric acid, citric acid, amino acids, such as glutamic acid or aspartic acid, maleic 35 acid.

acid, hydroxymaleic acid, methylmaleic acid, cyclohexanecarboxylic acid, adamantane-carboxylic acid, benzoic acid, salicylic acid, 4-aminosalicylic acid, phthalic acid, phenylacetic acid, mandelic acid, cinnamic acid, methane- or ethane-sulfonic acid, 2-hydroxyethanesulfonic acid, ethane-1,2-disulfonic acid, benzenesulfonic acid,
5 2-naphthalenesulfonic acid, 1,5-naphthalene-disulfonic acid, 2-, 3- or 4-methyl-benzenesulfonic acid, methylsulfuric acid, ethylsulfuric acid, dodecylsulfuric acid, N-cyclohexylsulfamic acid, N-methyl-, N-ethyl- or N-propyl-sulfamic acid, or other organic protonic acids, such as ascorbic acid.

10 For isolation or purification purposes it is also possible to use pharmaceutically unacceptable salts, for example picrates or perchlorates. For therapeutic use, only pharmaceutically acceptable salts or free compounds are employed (where applicable in the form of pharmaceutical preparations), and these are therefore preferred.

15 In view of the close relationship between the novel compounds in free form and those in the form of their salts, including those salts that can be used as intermediates, for example in the purification or identification of the novel compounds, any reference to the free compounds hereinbefore and hereinafter is to be understood as referring also to the corresponding salts, as appropriate and expedient.

20

The compound of the formula (I) may be administered in the form of a pro-drug which is broken down in the human or animal body to give a compound of the formula (I). Examples of pro-drugs include *in vivo* hydrolysable esters of a compound of the formula (I).

25

The compounds of formula (I) have valuable pharmacological properties. The invention also relates to compounds of formula (I) as defined hereinbefore for use as medicaments.

30

The efficacy of the compounds of the invention in inducing apoptosis in tumor cells can be demonstrated as follows:

35

Relative fluorescent activities of suitable tumor cell lines transfected with green fluorescent protein (GFP) are measured in the presence of compounds of the invention and of standard tumor drugs, using the method described in WO 99/35493. Suitable tumor cell lines are A20.2J, a BALB/c B cell lymphoma, PB-3c, an IL-3 dependent, non

tumorigenic mastocyte line isolated from the bone marrow of a DBA/2 mouse, Jurkat, a human acute T-cell leukemia cell line, K562, a human chronic myelogenous leukemia cell line, HL60, a human acute promyelocytic leukemia cell line, Ramos and Raji, human B-cell lymphoma cell lines, H9 and Hut78, human T-cell lymphoma cell lines, HeLa and KB, human squamous cell carcinoma cell lines, MCF7, SK-BR-3, PC3, HBL-100, SW480, H460 and H1792, human adenocarcinoma cell lines and HT-1080, a human fibrosarcoma cell line.

Preferred standard drugs as compounds for comparisons are: a) antimetabolites such as 5-fluorouracil (ICN), gemcitabine HCl (Gemzar™, Eli Lilly), b) alkylating agents such as oxaliplatin (Eloxantin™, Sanofi-Synthelabo), dacarbazine (Detimedac™, Medac), cyclophosphamide (Endoxan™, Asta) and carboplatin (Paraplatin™, Bristol-Meyers Squibb), c) cell-cycle inhibitor such as vinorelbine (Navelbine™, Robapharm), vinblastine (Velbe™, Eli Lilly), docetaxel (Taxotere™, Aventis), d) DNA breaker (topo-isomerase inhibitor, intercalator, strand breaker) such as doxorubicin HCl (Adriblastin™, Pharmacia-Upjohn), bleomycin (Asta-Medica), irinotecan (Campto™, Aventis), etoposide phosphate (Etopophos™, Bristol-Meyers Squibb), topotecan HCl, (Hycamtin™, GlaxoSmithKline), e) mixtures thereof, f) compounds interfering with the signal transduction pathway, such as caspase activity modifiers, agonists and antagonists of cell death receptors, modifiers of nucleases, phosphatases and kinases such as imatinib mesylate (Gleevec™, Novartis), dexamethasone, phorbol myristate acetate, cyclosporin A, quercetin, tamoxifen (Alexis Corporation, Switzerland).

Apoptosis is determined in a primary screen using a fluorescence plate reader and then in a secondary screen using FACS (fluorescence activated cell scanning). Compounds causing apoptosis without substantial cytotoxic side effects are chosen for further testing and characterization by using a combination of the following well established assays: A) Nuclear staining with Hoechst 33342 dye providing information about nuclear morphology and DNA fragmentation which are hallmarks of apoptosis. B) MTS proliferation assay measuring the metabolic activity of cells. Viable cells are metabolically active whereas cells with compromised respiratory chain show a reduced activity in this test. C) AnnexinV binding assay which reflects the phosphatidylserine content of the outer lipid bilayer of the plasma membrane. This event is considered an early hallmark of apoptosis. D) PI staining for cell cycle distribution which shows any alterations in the distribution among the different phases of the cell cycle. Cell cycle arresting points can be

determined. E) Proliferation assay monitoring DNA synthesis by incorporating bromodeoxyuridine (BrdU). Inhibitory effects on growth/proliferation can be directly determined. F) Cystein proteinase dependency, respectively caspase dependency are determined by using specific inhibitors. This provides information about possible involvement of specific proteases in the mechanisms.

On the basis of these studies, a compound of formula (I) according to the invention shows therapeutic efficacy especially against neoplastic diseases and autoimmune diseases. In particular, the compounds of the invention are active against malignancies, e.g. epithelial neoplasms, squamous cell neoplasms, basal cell neoplasms, transitional cell papillomas and carcinomas, adenomas und adenocarcinomas, adnexal and skin appendage neoplasms, mucoepidermoid neoplasms, cystic neoplasms, mucinous and serous neoplasms, ductal-, lobular and medullary neoplasms, acinar cell neoplasms, complex epithelial neoplasms, specialized gonadal neoplasms, paragangliomas and glomus tumors, naevi and melanomas, soft tissue tumors and sarcomas, fibromatous neoplasms, myxomatous neoplasms, lipomatous neoplasms, myomatous neoplasms, complex mixed and stromal neoplasms, fibroepithelial neoplasms, synovial like neoplasms, mesothelial neoplasms, germ cell neoplasms, trophoblastic neoplasms, mesonephromas, blood vessel tumors, lymphatic vessel tumors, osseous and chondromatous neoplasms, giant cell tumors, miscellaneous bone tumors, odontogenic tumors, gliomas, neuro-epitheliomatous neoplasms, meningiomas, nerve sheath tumors, granular cell tumors and alveolar soft part sarcomas, Hodgkin's and non Hodgkin's lymphomas, other lymphoreticular neoplasms, plasma cell tumors, mast cell tumors, immunoproliferative diseases, leukemias, miscellaneous myeloproliferative disorders, lymphoproliferative disorders and myelodysplastic syndromes.

The compounds of the invention are likewise active against autoimmune diseases, e.g. against systemic, discoid or subacute cutaneous lupus erythematosus, rheumatoid arthritis, antiphospholipid syndrome, CREST, progressive systemic sclerosis, mixed connective tissue disease (Sharp syndrome), Reiter's syndrome, juvenile arthritis, cold agglutinin disease, essential mixed cryoglobulinemia, rheumatic fever, ankylosing spondylitis, chronic polyarthritis, myasthenia gravis, multiple sclerosis, chronic inflammatory demyelinating polyneuropathy, Guillan-Barré syndrome, dermatomyositis/polymyositis, autoimmune hemolytic anemia, thrombocytopenic purpura, neutropenia, type I diabetes mellitus, thyroiditis (including Hashimoto's and Grave' disease), Addison's

- disease, polyglandular syndrome, pemphigus (vulgaris, foliaceus, sebaceous and vegetans), bullous and cicatricial pemphigoid, pemphigoid gestationis, epidermolysis bullosa acquisita, linear IgA disease, lichen sclerosus et atrophicus, morbus Duhring, psoriasis vulgaris, guttate, generalized pustular and localized pustular psoriasis, vitiligo,
- 5 alopecia areata, primary biliary cirrhosis, autoimmune hepatitis, all forms of glomerulonephritis, pulmonal hemorrhage (goodpasture syndrome), IgA nephropathy, pernicious anemia and autoimmune gastritis, inflammatory bowel diseases (including colitis ulcerosa and morbus Crohn), Behcet's disease, Celiac-Sprue disease, autoimmune uveitis, autoimmune myocarditis, granulomatous orchitis, aspermatogenesis without orchitis,
- 10 idiopathic and secondary pulmonary fibrosis, inflammatory diseases with a possibility of autoimmune pathogenesis, such as pyoderma gangrenosum, lichen ruber, sarcoidosis (including Löfgren and cutaneous/subcutaneous type), granuloma annulare, allergic type I and type IV immunological reaction, asthma bronchiale, pollinosis, atopic, contact and airborne dermatitis, large vessel vasculitis (giant cell and Takayasu's arteritis), medium
- 15 sized vessel vasculitis (polyarteritis nodosa, Kawasaki disease), small vessel vasculitis (Wegener's granulomatosis, Churg Strauss syndrome, microscopic polyangiitis, Henoch-Schoenlein purpura, essential cryoglobulinemic vasculitis, cutaneous leukoclastic angiitis), hypersensitivity syndromes, toxic epidermal necrolysis (Stevens-Johnson syndrome, erythema multiforme), diseases due to drug side effects, all forms of cutaneous, organ-
- 20 specific and systemic effects due to type I-VI (Coombs classification) immunologic forms of reaction, transplantation related pathologies, such as acute and chronic graft versus host and host versus graft disease, involving all organs (skin, heart, kidney, bone marrow, eye, liver, spleen, lung, muscle, central and peripheral nerve system, connective tissue, bone, blood and lymphatic vessel, genito-urinary system, ear, cartilage, primary and
- 25 secondary lymphatic system including bone marrow, lymph node, thymus, gastrointestinal tract, including oro-pharynx, esophageus, stomach, small intestine, colon, and rectum, including parts of above mentioned organs down to single cell level and substructures, e.g. stem cells).
- 30 A compound of formula (I) can be administered alone or in combination with one or more other therapeutic agents, possible combination therapy taking the form of fixed combinations, or the administration of a compound of the invention and one or more other therapeutic agents being staggered or given independently of one another, or the combined administration of fixed combinations and one or more other therapeutic agents.
- 35 A compound of formula (I) can, besides or in addition, be administered especially for

tumor therapy in combination with chemotherapy, radiotherapy, immunotherapy, surgical intervention, or a combination of these. Long-term therapy is equally possible as is adjuvant therapy in the context of other treatment strategies, as described above. Other possible treatments are therapy to maintain the patient's status after tumor regression, or even chemopreventive therapy, for example in patients at risk. Particularly preferred is the use of compounds of formula (I) in combination with radiotherapy.

Therapeutic agents for possible combination are especially one or more cytostatic or cytotoxic compounds, for example a chemotherapeutic agent or several selected from the group comprising indarubicin, cytarabine, interferon, hydroxyurea, bisulfan, or an inhibitor of polyamine biosynthesis, an inhibitor of protein kinase, especially of serine/threonine protein kinase, such as protein kinase C, or of tyrosine protein kinase, such as epidermal growth factor receptor tyrosine kinase, a cytokine, a negative growth regulator, such as TGF- β or IFN- β , an aromatase inhibitor, a classical cytostatic, an inhibitor of the interaction of an SH2 domain with a phosphorylated protein, an inhibitor of Bcl-2 and modulators of the Bcl-2 family members such as Bax, Bid, Bad, Bim, Nip3 and BH3-only proteins.

A compound according to the invention is not only for the (prophylactic and preferably therapeutic) management of humans, but also for the treatment of other warm-blooded animals, for example of commercially useful animals, for example rodents, such as mice, rabbits or rats, or guinea-pigs. Such a compound may also be used as a reference standard in the test systems described above to permit a comparison with other compounds.

With the groups of preferred compounds of formula (I) mentioned hereinafter, definitions of substituents from the general definitions mentioned hereinbefore may reasonably be used, for example, to replace more general definitions with more specific definitions or especially with definitions characterized as being preferred.

In particular, the invention refers to compounds of formula (I) wherein R represents aryl or heteroaryl optionally substituted by up to four substituents independently selected from alkyl, cycloalkyl, cycloalkyl-lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, lower alkoxy-lower alkoxy-lower alkyl, halo-lower alkoxy-lower alkyl, acyloxy-

- lower alkyl, heterocyclyl, heterocyclyl-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl, optionally substituted alkenyl, optionally substituted alkynyl, hydroxy, lower alkoxy, optionally substituted alkenyloxy, optionally substituted alkynyloxy, cycloalkoxy, halo-lower alkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, heterocycliloxy, heterocyclyl-lower alkoxy, optionally substituted phenyloxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy, optionally substituted heteroaryl-lower alkoxy, sulfamoyloxy, carbamoyloxy, lower alkylcarbonyloxy,
- 5 amino, monoalkylamino, dialkylamino, aminocarbonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylcarbonylamino wherein heterocyclyl is bound *via* a nitrogen atom, aminosulfonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylsulfonylamino wherein heterocyclyl
- 15 is bound *via* a nitrogen atom, lower alkoxycarbonylamino, lower alkylcarbonylamino wherein alkyl is optionally substituted by one or two substituents selected from optionally substituted phenyl, guanidyl, halogen, cyano, alkoxy, optionally substituted phenoxy, alkylmercapto and optionally substituted amino; lower alkenylcarbonylamino wherein alkenyl is optionally substituted by one or two substituents selected from lower alkyl, halo-
- 20 lower alkyl, optionally substituted phenyl, halogen, cyano, alkoxy and optionally substituted amino; amino-lower alkyl or amino-lower alkylamino, wherein the nitrogen atom is unsubstituted or substituted by one or two substituents selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted
- 25 heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl, lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclylcarbonyl, carboxy, lower alkoxycarbonyl, hydroxy-lower alkoxycarbonyl, lower alkoxy-lower
- 30 alkoxycarbonyl, optionally substituted phenyl-lower alkoxycarbonyl, cyano, lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl, halo-lower alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, and nitro; and wherein two adjacent substituents together with the atoms of aryl or heteroaryl may
- 35 form a 5 or 6 membered carbocyclic or heterocyclic ring;

X represents a bond or a group C=Y, wherein Y stands for oxygen;

- 5 R¹ and R², independently of each other, represent hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, optionally substituted arylalkyl, optionally substituted heteroarylalkyl, hydroxyalkyl, alkoxyalkyl, hydroxyalkoxyalkyl, alkoxyalkoxyalkyl, cyanoalkyl, optionally substituted alkenyl, optionally substituted alkynyl, or lower alkylcarbonyl wherein lower alkyl is optionally substituted by one or two substituents selected from aryl, optionally substituted amino, alkoxy and aryloxy;
- 10 R³, R⁴, R⁵ and R⁶, independently of each other, represent hydrogen, lower alkyl, halo-lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, lower alkoxy-lower alkoxy-lower alkyl, halo-lower alkoxy-lower alkyl, heterocyclyl, heterocyclyl-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl,
- 15 optionally substituted alkenyl, optionally substituted alkynyl, hydroxy, lower alkoxy, halo-lower alkoxy, cycloalkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, heterocyclioxy, heterocyclyl-lower alkoxy, optionally substituted phenyloxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy, optionally substituted heteroaryl-lower alkoxy,
- 20 amino, carbamoyl, sulfamoyl, amino-lower alkyl or amino-lower alkylamino, wherein in each case the nitrogen atom is unsubstituted or substituted by one or two substituents selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower
- 25 alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl, lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclylcarbonyl, carboxy, lower alkoxy carbonyl, hydroxy-lower alkoxy carbonyl, lower alkoxy-lower
- 30 alkoxy carbonyl, optionally substituted phenyl-lower alkoxy carbonyl, cyano, lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl, halo-lower alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, or nitro,

or R³ and R⁴, R⁴ and R⁵, or R⁵ and R⁶ together with the atoms of the phenyl ring form a 5 or 6 membered carbocyclic or heterocyclic ring;
and pharmaceutically acceptable salts thereof; for use as medicaments.

- 5 More particularly, the invention refers to compounds of formula (I) wherein
R represents phenyl, naphthyl, thienyl, furyl, thiazolyl, oxadiazolyl, thiadiazolyl, imidazolyl,
pyrazolyl, pyridinyl, pyrimidinyl, benzothienyl, benzofuryl, indolyl, benzisoxazolyl, each
optionally substituted by up to four substituents independently selected from
alkyl, cycloalkyl, cycloalkyl-lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-
10 lower alkyl, lower alkoxy-lower alkoxy-lower alkyl, halo-lower alkoxy-lower alkyl, acyloxy-
lower alkyl, heterocyclyl, heterocyclyl-lower alkyl, optionally substituted phenyl, optionally
substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted
heteroaryl-lower alkyl, optionally substituted alkenyl, optionally substituted alkynyl,
hydroxy, lower alkoxy, optionally substituted alkenyloxy, optionally substituted alkynyloxy,
15 cycloalkoxy, halo-lower alkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower
alkoxy-lower alkoxy, heterocyclyloxy, heterocyclyl-lower alkoxy, optionally substituted
phenyloxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy,
optionally substituted heteroaryl-lower alkoxy, sulfamoyloxy, carbamoyloxy, lower
alkylcarbonyloxy,
20 amino, monoalkylamino, dialkylamino, aminocarbonylamino wherein each of the two
amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl,
heterocyclylcarbonylamino wherein heterocyclyl is bound *via* a nitrogen atom,
aminosulfonylamino wherein each of the two amino groups is optionally substituted by
alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylsulfonylamino wherein heterocyclyl
25 is bound *via* a nitrogen atom, lower alkoxy carbonylamino, lower alkylcarbonylamino
wherein alkyl is optionally substituted by one or two substituents selected from optionally
substituted phenyl, guanidyl, halogen, cyano, alkoxy, optionally substituted phenoxy,
alkylmercapto and optionally substituted amino; lower alkenylcarbonylamino wherein
alkenyl is optionally substituted by one or two substituents selected from lower alkyl, halo-
30 lower alkyl, optionally substituted phenyl, halogen, cyano, alkoxy and optionally
substituted amino; amino-lower alkyl or amino-lower alkylamino, wherein the nitrogen
atom is unsubstituted or substituted by one or two substituents selected from lower alkyl,
cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally
substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted

heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl, lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclylcarbonyl,

5 lower alkylsulfinyl, halo-lower alkylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, halogen, and nitro;

and wherein two adjacent substituents together with the atoms of aryl or heteroaryl may form a 5 or 6 membered carbocyclic or heterocyclic ring;

10 X represents a bond or a group C=Y, wherein Y stands for oxygen;

R¹ and R², independently of each other, represent hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, optionally substituted arylalkyl, optionally substituted heteroarylalkyl, hydroxyalkyl, alkoxyalkyl, hydroxyalkoxyalkyl, alkoxyalkoxyalkyl, optionally substituted alkenyl,

15 optionally substituted alkynyl, lower alkylcarbonyl wherein lower alkyl is optionally substituted by one or two substituents selected from aryl, optionally substituted amino or aroyl, or optionally substituted phenylcarbonyl; or R¹ and R² together with the atom they are bound to form heterocyclyl;

20 R³, R⁴, R⁵ and R⁶, independently of each other, represent hydrogen, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, cycloalkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, heterocycliloxy, heterocyclyl-lower alkoxy, optionally substituted phenyloxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy, optionally substituted heteroaryl-lower alkoxy,

25 amino, carbamoyl, sulfamoyl, amino-lower alkyl or amino-lower alkylamino, wherein in each case the nitrogen atom is unsubstituted or substituted by one or two substituents selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl,

30 lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclylcarbonyl,

35 carboxy, lower alkoxy carbonyl, hydroxy-lower alkoxy carbonyl, lower alkoxy-lower alkoxy carbonyl, optionally substituted phenyl-lower alkoxy carbonyl, cyano,

lower alkylsulfinyl, halo-lower alkylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, halogen, or nitro;

or R³ and R⁴, R⁴ and R⁵, or R⁵ and R⁶ together represent methylenedioxy;

and pharmaceutically acceptable salts thereof, for use as a medicament.

5

Preferably, the invention refers to compounds of formula (I) wherein

R represents phenyl, naphthyl, thienyl, furyl, thiazolyl, oxadiazolyl, thiadiazolyl, imidazolyl, pyrazolyl, pyridinyl, pyrimidinyl, benzothienyl, benzofuryl, indolyl, benzisoxazolyl, optionally substituted by up to four substituents independently selected from

10 alkyl, halo-lower alkyl, phenyl, optionally substituted heteroaryl, lower alkoxy, optionally substituted alkenyloxy, optionally substituted alkynyloxy, lower alkoxy-lower alkoxy, amino, monoalkylamino, dialkylamino, aminocarbonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylcarbonylamino wherein heterocyclyl is bound via a nitrogen atom,

15 aminosulfonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylsulfonylamino wherein heterocyclyl is bound via a nitrogen atom, lower alkoxycarbonylamino, lower alkylcarbonylamino wherein alkyl is optionally substituted by alkoxy or optionally substituted amino; lower alkenylcarbonylamino wherein alkenyl is optionally substituted by alkoxy or optionally substituted amino; lower alkylsulfinyl, halo-lower alkylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl and halogen;

and wherein two adjacent substituents together with the atoms of aryl or heteroaryl may form a 5 or 6 membered carbocyclic or heterocyclic ring;

X represents a bond or a group C=Y, wherein Y stands for oxygen;

25 R¹ and R², independently of each other, represent hydrogen or lower alkylcarbonyl; R³, R⁴, R⁵ and R⁶, independently of each other, represent hydrogen, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy, carboxy, lower alkoxycarbonyl, cyano or halogen; and pharmaceutically acceptable salts thereof, for use as a medicament.

30 Most preferably, the invention relates to the compounds of the Examples and pharmaceutically acceptable salts thereof for use as a medicament, especially to the compounds of Examples 1, 5, 8, 9, 12, 13, 14, 15, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 43, 44, and 45, and to pharmaceutically acceptable salts thereof.

Furthermore, the invention refers to novel compounds of formula (I) as such, in particular to compounds of formula (I) wherein

R represents aryl or heteroaryl optionally substituted by up to four substituents

5 independently selected from

alkyl, cycloalkyl, cycloalkyl-lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, lower alkoxy-lower alkoxy-lower alkyl, halo-lower alkoxy-lower alkyl, acyloxy-lower alkyl, heterocyclyl, heterocyclyl-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl, optionally substituted alkenyl, optionally substituted alkynyl, hydroxy, lower alkoxy, optionally substituted alkenyloxy, optionally substituted alkynyloxy, cycloalkoxy, halo-lower alkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, heterocycliloxy, heterocyclyl-lower alkoxy, optionally substituted phenyloxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy, optionally substituted heteroaryl-lower alkoxy, sulfamoyloxy, carbamoyloxy, lower alkylcarbonyloxy,

amino, monoalkylamino, dialkylamino, aminocarbonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylcarbonylamino wherein heterocyclyl is bound *via* a nitrogen atom,

20 aminosulfonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylsulfonylamino wherein heterocyclyl is bound *via* a nitrogen atom, lower alkoxycarbonylamino, lower alkylcarbonylamino wherein alkyl is optionally substituted by one or two substituents selected from optionally substituted phenyl, guanidyl, halogen, cyano, alkoxy, optionally substituted phenoxy, alkylmercapto and optionally substituted amino; lower alkenylcarbonylamino wherein alkenyl is optionally substituted by one or two substituents selected from lower alkyl, halo-lower alkyl, optionally substituted phenyl, halogen, cyano, alkoxy and optionally substituted amino; amino-lower alkyl or amino-lower alkylamino, wherein the nitrogen atom is unsubstituted or substituted by one or two substituents selected from lower alkyl, 30 cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl, lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclylcarbonyl,

carboxy, lower alkoxycarbonyl, hydroxy-lower alkoxycarbonyl, lower alkoxy-lower alkoxycarbonyl, optionally substituted phenyl-lower alkoxycarbonyl, cyano, lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl, halo-lower alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, and nitro; and wherein two adjacent substituents together with the atoms of aryl or heteroaryl may form a 5 or 6 membered carbocyclic or heterocyclic ring;

X represents oxygen; a group C=Y, wherein Y stands for oxygen, nitrogen substituted by hydroxy, alkoxy or optionally substituted amino; an ethylenediyl group optionally substituted by alkyl; or a group CR⁷R⁸;

R¹ and R², independently of each other, represent hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, optionally substituted arylalkyl, optionally substituted heteroarylalkyl, hydroxyalkyl, alkoxyalkyl, hydroxyalkoxyalkyl, alkoxyalkoxyalkyl, cyanoalkyl, optionally substituted alkenyl, optionally substituted alkynyl, or lower alkylcarbonyl wherein lower alkyl is optionally substituted by one or two substituents selected from aryl, optionally substituted amino, alkoxy and aryloxy,

R³, R⁴, R⁵ and R⁶, independently of each other, represent hydrogen, lower alkyl, halo-lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, lower alkoxy-lower alkoxy-lower alkyl, halo-lower alkoxy-lower alkyl, heterocyclyl, heterocyclyl-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl, optionally substituted alkenyl, optionally substituted alkynyl, hydroxy, lower alkoxy, halo-lower alkoxy, cycloalkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, heterocyclyloxy, heterocyclyl-lower alkoxy, optionally substituted phenyloxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy, optionally substituted heteroaryl-lower alkoxy, amino, carbamoyl, sulfamoyl, amino-lower alkyl or amino-lower alkylamino, wherein in each case the nitrogen atom is unsubstituted or substituted by one or two substituents selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower

alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl,

lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclylcarbonyl,

5 carboxy, lower alkoxy carbonyl, hydroxy-lower alkoxy carbonyl, lower alkoxy-lower alkoxy carbonyl, optionally substituted phenyl-lower alkoxy carbonyl, cyano, lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl, halo-lower alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, or nitro,

10

or R³ and R⁴, R⁴ and R⁵, or R⁵ and R⁶ together with the atoms of the phenyl ring form a 5 or 6 membered carbocyclic or heterocyclic ring;

R⁷ represents hydrogen, lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, lower alkenyl, lower
15 alkynyl, optionally substituted phenyl, lower alkoxy, lower alkenyloxy, lower alkynyloxy;
R⁸ represents hydrogen, lower alkyl, hydroxy, lower alkoxy or lower alkenyloxy, or
R⁷ and R⁸ together with the carbon they are bound to form a 5 or 6 membered carbocyclic
or heterocyclic ring;
and salts thereof.

20

More particularly, the invention refers to compounds of formula (I) wherein

R represents phenyl, naphthyl, thienyl, furyl, thiazolyl, oxadiazolyl, thiadiazolyl, imidazolyl,
pyrazolyl, pyridinyl, pyrimidinyl, benzothienyl, benzofuryl, indolyl, benzoisoxazolyl,
optionally substituted by up to four substituents independently selected from

25 alkyl, cycloalkyl, cycloalkyl-lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-
lower alkyl, lower alkoxy-lower alkoxy-lower alkyl, halo-lower alkoxy-lower alkyl, acyloxy-
lower alkyl, heterocyclyl, heterocyclyl-lower alkyl, optionally substituted phenyl, optionally
substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted
heteroaryl-lower alkyl, optionally substituted alkenyl, optionally substituted alkynyl,
30 hydroxy, lower alkoxy, optionally substituted alkenyloxy, optionally substituted alkynyloxy,
cycloalkoxy, halo-lower alkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower
alkoxy-lower alkoxy, heterocycliloxy, heterocyclyl-lower alkoxy, optionally substituted
phenyloxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy,
optionally substituted heteroaryl-lower alkoxy, sulfamoyloxy, carbamoyloxy, lower
35 alkylcarbonyloxy,

amino, monoalkylamino, dialkylamino, aminocarbonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylcarbonylamino wherein heterocyclyl is bound *via* a nitrogen atom, aminosulfonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylsulfonylamino wherein heterocyclyl is bound *via* a nitrogen atom, lower alkoxy-carbonylamino, lower alkylcarbonylamino wherein alkyl is optionally substituted by one or two substituents selected from optionally substituted phenyl, guanidyl, halogen, cyano, alkoxy, optionally substituted phenoxy, alkylmercapto and optionally substituted amino; lower alkenylcarbonylamino wherein alkenyl is optionally substituted by one or two substituents selected from lower alkyl, halo-lower alkyl, optionally substituted phenyl, halogen, cyano, alkoxy and optionally substituted amino; amino-lower alkyl or amino-lower alkylamino, wherein the nitrogen atom is unsubstituted or substituted by one or two substituents selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl, lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclylcarbonyl, carboxy, lower alkoxy-carbonyl, hydroxy-lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, optionally substituted phenyl-lower alkoxy-carbonyl, cyano, lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl, halo-lower alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, and nitro; and wherein two adjacent substituents together with the atoms of aryl or heteroaryl may form a 5 or 6 membered carbocyclic or heterocyclic ring;

X represents oxygen; a group C=Y, wherein Y stands for oxygen, nitrogen substituted by hydroxy, alkoxy or optionally substituted amino; an ethylenediyl group optionally substituted by alkyl; or a group CR⁷R⁸;

R¹ and R², independently of each other, represent hydrogen, lower alkylcarbonyl or optionally substituted phenylcarbonyl;

R^3 , R^4 , R^5 and R^6 , independently of each other, represent hydrogen, lower alkyl, halo-lower alkyl, alkoxy, hydroxy-lower alkoxy, lower-alkoxy-lower-alkoxy; amino, carbamoyl, sulfamoyl, amino-lower alkyl or amino-lower alkylamino, wherein in each case the nitrogen atom is unsubstituted or substituted by one or two substituents selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylaminocarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl, carboxy, lower alkoxycarbonyl, hydroxy-lower alkoxycarbonyl, lower alkoxy-lower alkoxycarbonyl, optionally substituted phenyl-lower alkoxycarbonyl, cyano, lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl, halo-lower alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, or nitro,

or R^3 and R^4 , R^4 and R^5 , or R^5 and R^6 together represent methylenedioxy;

R^7 represents hydrogen, lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, lower alkenyl, lower alkynyl, optionally substituted phenyl, hydroxy, lower alkoxy, lower alkenyloxy, lower alkynyloxy;

R^8 represents hydrogen, lower alkyl, lower alkoxy or lower alkenyloxy, or R^7 and R^8 together with the carbon they are bound to form a 5 or 6 membered carbocyclic or heterocyclic ring; and salts thereof.

Preferably, the invention refers to compounds of formula (I) wherein R represents phenyl, naphthyl, thienyl, furyl, thiazolyl, oxadiazolyl, thiadiazolyl, imidazolyl, pyrazolyl, pyridinyl, pyrimidinyl, benzothienyl, benzofuryl, indolyl, benzoisoxazolyl, optionally substituted by up to four substituents independently selected from alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, lower alkoxy-lower alkoxy-lower alkyl, halo-lower alkoxy-lower alkyl, acyloxy-lower alkyl, heterocyclyl, heterocyclyl-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl, optionally substituted alkenyl, optionally substituted alkynyl, hydroxy, lower alkoxy, optionally substituted alkenyloxy, optionally substituted alkynyloxy, cycloalkoxy, halo-lower alkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower

alkoxy-lower alkoxy, heterocyclyloxy, heterocyclyl-lower alkoxy, optionally substituted phenyloxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy, optionally substituted heteroaryl-lower alkoxy, sulfamoyloxy, carbamoyloxy, lower alkylcarbonyloxy,

- 5 amino, monoalkylamino, dialkylamino, aminocarbonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylcarbonylamino wherein heterocyclyl is bound *via* a nitrogen atom, aminosulfonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylsulfonylamino wherein heterocyclyl
- 10 is bound *via* a nitrogen atom, lower alkoxycarbonylamino, lower alkylcarbonylamino wherein alkyl is optionally substituted by one or two substituents selected from optionally substituted phenyl, guanidyl, halogen, cyano, alkoxy, optionally substituted phenoxy, alkylmercapto and optionally substituted amino; lower alkenylcarbonylamino wherein alkenyl is optionally substituted by one or two substituents selected from lower alkyl, halo-
- 15 lower alkyl, optionally substituted phenyl, halogen, cyano, alkoxy and optionally substituted amino; amino-lower alkyl or amino-lower alkylamino, wherein the nitrogen atom is unsubstituted or substituted by one or two substituents selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted
- 20 heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl, lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclylcarbonyl, carboxy, lower alkoxycarbonyl, hydroxy-lower alkoxycarbonyl, lower alkoxy-lower
- 25 alkoxycarbonyl, optionally substituted phenyl-lower alkoxycarbonyl, cyano, lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl, halo-lower alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, and nitro; and wherein two adjacent substituents together with the atoms of aryl or heteroaryl may
- 30 form a 5 or 6 membered carbocyclic or heterocyclic ring;

X represents a group C=Y, wherein Y stands for oxygen, nitrogen substituted by hydroxy, alkoxy or optionally substituted amino; or a group CR⁷R⁸;

- 35 R¹ and R², independently of each other, represent hydrogen or lower alkylcarbonyl;

- R³, R⁴, R⁵ and R⁶, independently of each other, represent hydrogen, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy, carboxy, lower alkoxycarbonyl, cyano, halogen or nitro; R⁷ represents hydrogen, lower alkyl, hydroxy or lower alkoxy; R⁸ represents hydrogen, lower alkyl or lower alkoxy; or
- 5 R⁷ and R⁸ together represent 1,2-ethylenedioxy or 3-alkoxypropylene-1,2-dioxy; and salts thereof.

Alternatively, the invention relates to compound of formula (I)

- 10 R represents heteroaryl optionally substituted by up to four substituents independently selected from
- alkyl, cycloalkyl, cycloalkyl-lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, lower alkoxy-lower alkoxy-lower alkyl, halo-lower alkoxy-lower alkyl, acyloxy-lower alkyl, heterocyclyl, heterocyclyl-lower alkyl, optionally substituted phenyl, optionally
- 15 substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl, optionally substituted alkenyl, optionally substituted alkynyl, hydroxy, lower alkoxy, optionally substituted alkenyloxy, optionally substituted alkynyloxy, cycloalkoxy, halo-lower alkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, heterocyclyloxy, heterocyclyl-lower alkoxy, optionally substituted
- 20 phenyloxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy, optionally substituted heteroaryl-lower alkoxy, sulfamoyloxy, carbamoyloxy, lower alkylcarbonyloxy,
- amino, monoalkylamino, dialkylamino, aminocarbonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl,
- 25 heterocyclylcarbonylamino wherein heterocyclyl is bound *via* a nitrogen atom, aminosulfonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylsulfonylamino wherein heterocyclyl is bound *via* a nitrogen atom, lower alkoxycarbonylamino, lower alkylcarbonylamino wherein alkyl is optionally substituted by one or two substituents selected from optionally
- 30 substituted phenyl, guanidyl, halogen, cyano, alkoxy, optionally substituted phenoxy, alkylmercapto and optionally substituted amino; lower alkenylcarbonylamino wherein alkenyl is optionally substituted by one or two substituents selected from lower alkyl, halo-lower alkyl, optionally substituted phenyl, halogen, cyano, alkoxy and optionally substituted amino; amino-lower alkyl or amino-lower alkylamino, wherein the nitrogen
- 35 atom is unsubstituted or substituted by one or two substituents selected from lower alkyl,

- cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl,
- 5 lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclylcarbonyl, carboxy, lower alkoxy carbonyl, hydroxy-lower alkoxy carbonyl, lower alkoxy-lower alkoxy carbonyl, optionally substituted phenyl-lower alkoxy carbonyl, cyano, lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl, halo-lower
- 10 alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, and nitro; and wherein two adjacent substituents together with the atoms of aryl or heteroaryl may form a 5 or 6 membered carbocyclic or heterocyclic ring;
- 15 X represents a bond;
- R^1 and R^2 , independently of each other, represent hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, optionally substituted arylalkyl, optionally substituted heteroarylalkyl, hydroxyalkyl, alkoxyalkyl, hydroxyalkoxyalkyl, alkoxyalkoxyalkyl, cyanoalkyl, optionally substituted
- 20 alkenyl, optionally substituted alkynyl, or lower alkylcarbonyl wherein lower alkyl is optionally substituted by one or two substituents selected from aryl, optionally substituted amino, alkoxy and aryloxy,
- R^3 , R^4 , R^5 and R^6 , independently of each other, represent hydrogen, lower alkyl, halo-
- 25 lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, lower alkoxy-lower alkoxy-lower alkyl, halo-lower alkoxy-lower alkyl, heterocyclyl, heterocyclyl-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl, optionally substituted alkenyl, optionally substituted alkynyl,
- 30 hydroxy, lower alkoxy, halo-lower alkoxy, cycloalkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, heterocycliloxy, heterocyclyl-lower alkoxy, optionally substituted phenyloxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy, optionally substituted heteroaryl-lower alkoxy, amino, carbamoyl, sulfamoyl, amino-lower alkyl or amino-lower alkylamino, wherein in
- 35 each case the nitrogen atom is unsubstituted or substituted by one or two substituents

selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl; optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl,

5 lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclylcarbonyl, carboxy, lower alkoxy carbonyl, hydroxy-lower alkoxy carbonyl, lower alkoxy-lower alkoxy carbonyl, optionally substituted phenyl-lower alkoxy carbonyl, cyano,

10 lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl, halo-lower alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, or nitro,

or R³ and R⁴, R⁴ and R⁵, or R⁵ and R⁶ together with the atoms of the phenyl ring form a 5 or 6 membered carbocyclic or heterocyclic ring;

15 and salts thereof;

with the exception of compounds of formula (I) wherein R represents 4-pyridyl, 3,5-diaminotriazinyl, 3-amino-5-piperidinotriazinyl, 5-methyl-1,3,4-oxadiazol-2-yl, 5-amino-1,3,4-thiadiazol-2-yl and 1H-2-tetrazolyl, X is a bond, R¹ is hydrogen or acetyl, and R², R³, R⁴, R⁵ and R⁶ are all hydrogen.

20

More particularly, the invention refers to compounds of formula (I) wherein R represents thienyl, furyl, thiazolyl, oxadiazolyl, thiadiazolyl, imidazolyl, pyrazolyl, pyridinyl, pyrimidinyl, benzothienyl, benzofuryl, indolyl, benzoisoxazolyl, optionally substituted by up to four substituents independently selected from alkyl, cycloalkyl, cycloalkyl-lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, lower alkoxy-lower alkoxy-lower alkyl, halo-lower alkoxy-lower alkyl, acyloxy-lower alkyl, heterocyclyl, heterocyclyl-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl, optionally substituted alkenyl, optionally substituted alkynyl, hydroxy, lower alkoxy, optionally substituted alkenyloxy, optionally substituted alkynyloxy, cycloalkoxy, halo-lower alkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, heterocyclyloxy, heterocyclyl-lower alkoxy, optionally substituted phenyloxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy,

25

30

35

- optionally substituted heteroaryl-lower alkoxy, sulfamoyloxy, carbamoyloxy, lower alkylcarbonyloxy,
- 5 amino, monoalkylamino, dialkylamino, aminocarbonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylcarbonylamino wherein heterocyclyl is bound *via* a nitrogen atom, aminosulfonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylsulfonylamino wherein heterocyclyl is bound *via* a nitrogen atom, lower alkoxycarbonylamino, lower alkylcarbonylamino wherein alkyl is optionally substituted by one or two substituents selected from optionally
- 10 substituted phenyl, guanidyl, halogen, cyano, alkoxy, optionally substituted phenoxy, alkylmercapto and optionally substituted amino; lower alkenylcarbonylamino wherein alkenyl is optionally substituted by one or two substituents selected from lower alkyl, halo-lower alkyl, optionally substituted phenyl, halogen, cyano, alkoxy and optionally substituted-amino; amino-lower alkyl or amino-lower-alkylamino; wherein the nitrogen
- 15 atom is unsubstituted or substituted by one or two substituents selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl,
- 20 lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclylcarbonyl, carboxy, lower alkoxycarbonyl, hydroxy-lower alkoxycarbonyl, lower alkoxy-lower alkoxycarbonyl, optionally substituted phenyl-lower alkoxycarbonyl, cyano, lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl, halo-lower
- 25 alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, and nitro; and wherein two adjacent substituents together with the atoms of aryl or heteroaryl may form a 5 or 6 membered carbocyclic or heterocyclic ring;

- 30 X represents a bond;

R¹ and R², independently of each other, represent hydrogen, lower alkylcarbonyl or optionally substituted phenylcarbonyl;

R^3 , R^4 , R^5 and R^6 , independently of each other, represent hydrogen, lower alkyl, halo-lower-alkyl, alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower-alkoxy, amino, carbamoyl, sulfamoyl, amino-lower alkyl or amino-lower alkylamino, wherein in each case the nitrogen atom is unsubstituted or substituted by one or two substituents selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylaminocarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl, carboxy, lower alkoxycarbonyl, hydroxy-lower alkoxycarbonyl, lower alkoxy-lower alkoxycarbonyl, optionally substituted phenyl-lower alkoxycarbonyl, cyano, lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl, halo-lower alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, or nitro,

or R^3 and R^4 , R^4 and R^5 , or R^5 and R^6 together represent methylenedioxy; and salts thereof;

with the exception of compounds of formula (I) wherein R represents 4-pyridyl, 5-methyl-1,3,4-oxadiazol-2-yl and 5-amino-1,3,4-thiadiazol-2-yl, X is a bond, R^1 is hydrogen or acetyl, and R^2 , R^3 , R^4 , R^5 and R^6 are all hydrogen.

Preferably, the invention refers to compounds of formula (I) wherein R represents thienyl, furyl, thiazolyl, oxadiazolyl, thiadiazolyl, imidazolyl, pyrazolyl, pyridinyl, pyrimidinyl, benzothienyl, benzofuryl, indolyl, benzoisoxazolyl, optionally substituted by up to four substituents independently selected from alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, lower alkoxy-lower alkoxy-lower alkyl, halo-lower alkoxy-lower alkyl, acyloxy-lower alkyl, heterocyclyl, heterocyclyl-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl, optionally substituted alkenyl, optionally substituted alkynyl, hydroxy, lower alkoxy, optionally substituted alkenyloxy, optionally substituted alkynyloxy, cycloalkoxy, halo-lower alkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, heterocyclyloxy, heterocyclyl-lower alkoxy, optionally substituted phenyloxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy,

optionally substituted heteroaryl-lower alkoxy, sulfamoyloxy, carbamoyloxy, lower alkylcarbonyloxy,
amino, monoalkylamino, dialkylamino, aminocarbonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl,
5 heterocyclylcarbonylamino wherein heterocyclyl is bound *via* a nitrogen atom, aminosulfonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylsulfonylamino wherein heterocyclyl is bound *via* a nitrogen atom, lower alkoxycarbonylamino, lower alkylcarbonylamino wherein alkyl is optionally substituted by one or two substituents selected from optionally
10 substituted phenyl, guanidyl, halogen, cyano, alkoxy, optionally substituted phenoxy, alkylmercapto and optionally substituted amino; lower alkenylcarbonylamino wherein alkenyl is optionally substituted by one or two substituents selected from lower alkyl, halo-lower alkyl, optionally substituted phenyl, halogen, cyano, alkoxy and optionally substituted amino; amino-lower alkyl or amino-lower alkylamino, wherein the nitrogen
15 atom is unsubstituted or substituted by one or two substituents selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl,
20 lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclylcarbonyl, carboxy, lower alkoxycarbonyl, hydroxy-lower alkoxycarbonyl, lower alkoxy-lower alkoxycarbonyl, optionally substituted phenyl-lower alkoxycarbonyl, cyano, lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl, halo-lower
25 alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, and nitro; and wherein two adjacent substituents together with the atoms of aryl or heteroaryl may form a 5 or 6 membered carbocyclic or heterocyclic ring;

30 X represents a bond;

R¹ and R², independently of each other, represent hydrogen or lower alkylcarbonyl;

R³, R⁴, R⁵ and R⁶, independently of each other, represent hydrogen, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy, carboxy, lower alkoxycarbonyl, cyano, halogen or nitro;
35

and salts thereof;

with the exception of compounds of formula (I) wherein R represents 4-pyridyl, 5-methyl-
1,3,4-oxadiazol-2-yl and 5-amino-1,3,4-thiadiazol-2-yl, X is a bond, R¹ is hydrogen or
5 acetyl, and R², R³, R⁴, R⁵ and R⁶ are all hydrogen.

Especially, the invention relates to the use of a compound of formula (I), a prodrug or a
pharmaceutically acceptable salt of such a compound for the preparation of a
pharmaceutical composition for the treatment of a neoplastic disease, autoimmune
10 disease, transplantation related pathology and/or degenerative disease.

Furthermore, the invention provides a method for the treatment of a neoplastic disease,
autoimmune disease, transplantation related pathology and/or degenerative disease,
which comprises administering a compound of formula (I), a prodrug or a
15 pharmaceutically acceptable salt thereof, wherein the radicals and symbols have the
meanings as defined above, in a quantity effective against said disease, to a warm-
blooded animal requiring such treatment.


Method of preparation

20

A compound of the invention may be prepared by processes that, though not applied
hitherto for the new compounds of the present invention, are known *per se*, in particular

A) a process, wherein a compound of formula (II)

25



wherein R¹, R², R³, R⁴, R⁵ and R⁶ are defined as for formula (I), or a derivative thereof with
functional groups in protected form and/or a salt thereof, is alkylated with a halide of the
formula (III)


30




wherein R is as defined for formula (I) and Z is a nucleophilic leaving group;

5 or

B) a process, wherein a compound of formula (IV)



10 wherein R¹ and R² are defined as for formula (I) and R' is alkyl, or a derivative thereof with functional groups in protected form and/or a salt thereof, is reacted with an ortho-phenylene diamine of formula (V),



15 wherein R, R³, R⁴, R⁵ and R⁶ are defined as for formula (I), or a derivative thereof wherein functional groups are in protected form and/or a salt thereof;

any protecting groups in a protected derivative of a compound of the formula (I) are removed;

20 and, if so desired, an obtainable compound of formula (I) is converted into another compound of formula (I), a free compound of formula (I) is converted into a salt, an obtainable salt of a compound of formula (I) is converted into the free compound or another salt, and/or a mixture of isomeric compounds of formula (I) is separated into the individual isomers.

Suitable nucleophilic leaving groups Z in an alkylating agent of formula (III) are for example halides, e.g. chloride, bromide or iodide, or sulfonates, e.g. aromatic sulfonic acid esters such as benzenesulfonates, p-toluenesulfonates or p-nitrobenzenesulfonates, or also methanesulfonate or trifluoromethanesulfonate. Also other customary leaving groups are considered, e.g. ammonium salts, azides, diazonium salts, di(p-toluenesulfonyl)-amines, nitrates, oxonium salts, sulfonium salts, or phosphonium salts.

Alkylation of a compound of formula (II) with an alkylating agent of formula (III) is performed in a manner known *per se*, usually in the presence of a suitable polar or dipolar aprotic solvent, with cooling or heating, for example in a temperature range from approximately -30°C to approximately +150°C, especially approximately around 0°C to room temperature. Optionally a suitable base is added, in particular a tertiary amine base such as triethylamine or diisopropylethylamine, or an inorganic basic salt, e.g. potassium or sodium carbonate.

Cyclization of an imino ether (IV) with a phenylene diamine (V) is performed in a manner known *per se*, usually in the presence of a suitable polar or dipolar aprotic solvent, preferably in an alcohol, with cooling or heating, for example in a temperature range from approximately 30°C to approximately +150°C, under conditions taking into account the critical stability of the phenylene diamine (V).

If one or more other functional groups, for example carboxy, hydroxy or amino, are or need to be protected in a compound of formula (II), (III), (IV) or (V), because they should not take part in the reaction, these are such protecting groups as are usually applied in the synthesis of amides, in particular peptide compounds, cephalosporins, penicillins, nucleic acid derivatives and sugars.

The protecting groups may already be present in precursors and should protect the functional groups concerned against unwanted secondary reactions, such as alkylations, acylations, etherifications, esterifications, oxidations, solvolysis, and similar reactions. It is a characteristic of protecting groups that they lend themselves readily, i.e. without undesired secondary reactions, to removal, typically by solvolysis, reduction, photolysis or also by enzyme activity, for example under conditions analogous to physiological conditions, and that they are not present in the end products. The specialist knows, or can

easily establish, which protecting groups are suitable with the reactions mentioned hereinabove and hereinafter:

5 The protection of such functional groups by such protecting groups, the protecting groups themselves, and their removal reactions are described for example in standard reference books for peptide synthesis and in special books on protective groups such as J. F. W. McOmie, "Protective Groups in Organic Chemistry", Plenum Press, London and New York 1973, in "Methoden der organischen Chemie" (Methods of organic chemistry), Houben-Weyl, 4th edition, Volume 15/I, Georg Thieme Verlag, Stuttgart 1974, and in
10 T. W. Greene, "Protective Groups in Organic Synthesis", Wiley, New York.

Other processes may be considered for the preparation of compounds of formula (I) wherein X is a bond and R is heteroaryl. For example, a compound of formula (I) wherein X is a bond and R is $-C(=O)NHNH_2$, i.e. a hydrazide, may be transformed by reaction with
15 an orthoester, and amidine or an acylating agent followed by dehydration to give a corresponding compound wherein R is a 5-substituted 1,3,4-oxadiazol-2-yl group. When R is $-C(NH_2)=N-OH$, i.e. a hydroxamic acid amide, the corresponding reaction leads to a group R being a 5-substituted 1,2,4-oxadiazol-3-yl function. When R is $-C(NH_2)=NH$, i.e. an amidine, reaction with a 1,3-diketone gives a 4,6-disubstituted pyrimidin-2-yl function.
20 Other heterocycles may be formed in analogous reactions.

In the additional process steps, carried out as desired, functional groups of the starting compounds which should not take part in the reaction may be present in unprotected form or may be protected for example by one or more of the protecting groups mentioned
25 hereinabove under "protecting groups". The protecting groups are then wholly or partly removed according to one of the methods described there.

In the conversion of an obtainable compound of formula (I) into another compound of formula (I), X with the meaning $C=Y$ wherein Y is oxygen may, for example, be reduced to
30 X with the meaning CR^7R^8 wherein R^7 is hydrogen or hydroxy and R^8 is hydrogen. Suitable reducing agents are known in the art, and are, for example, metal hydrides, e.g. $LiAlH_4$, $LiAl(OCH_3)_3H$ or other alkoxy-substituted lithium hydrides, $NaBH_4$, or BH_3 , optionally in the presence of a Lewis base, e.g. $AlCl_3$ or BF_3 , or also with catalytical hydrogenation with hydrogen and a suitable noble metal catalyst. Through the choice of
35 catalyst and reaction conditions, it can be influenced whether the reaction stops at the

alcohol stage (R^7 hydroxy) or to the fully saturated methylene stage (R^7 hydrogen).

- A compound of formula (I) wherein X is $C=Y$ and Y is oxygen may be reacted with an optionally O-substituted hydroxylamine to give the corresponding oxime or oxime ether of formula (I) wherein X is $C=Y$ and Y is nitrogen substituted by hydroxy or alkoxy. By reaction with an optionally substituted hydrazine, the corresponding hydrazone of formula (I) wherein X is $C=Y$ and Y is nitrogen substituted by optionally substituted amino is formed.
- 10 A compound of formula (I) wherein X is $C=Y$ and Y is oxygen may be reacted with a suitably substituted alcohol to give the corresponding acetal, i.e. a compound of formula (I) wherein X is CR^7R^8 and R^7 and R^8 represent alkoxy, in the presence of an acid catalyst and optionally a water binding agent and/or a water trap. Compounds of formula (I) wherein R^7 and R^8 are part of a 1,3-dioxolane or a 1,3-dioxane may be obtained analogously by reaction of a compound of formula (I) wherein X is $C=O$ and Y is oxygen with a polyalcohol, such as glycol, propane-1,3-diol, glycerol and the like.

- An obtainable compound of formula (I), wherein R^1 and/or R^2 is hydrogen, may be alkylated or acylated with a compound of formula R^1-Z or R^2-Z , respectively, wherein Z is a nucleophilic leaving group as described above, to give a compound of formula (I), wherein R^1 and/or R^2 is different from hydrogen. Preferred acylation conditions include the use of acid anhydrides and acid chlorides at elevated temperatures, typically in a range from approximately $+30^\circ\text{C}$ to approximately $+150^\circ\text{C}$. An acidic or basic catalyst may be employed if desired. A compound of formula (I) wherein R^1 and/or R^2 is alkyl may be obtained by alkylation of the parent compound of formula (I). Typical reaction conditions allowing this transformation include the combination of a strong base, such as a metal hydride or a metal alcoholate and a compound of formula R^1-Z or R^2-Z .

- Further amino groups present in an aryl or heteroaryl group R or in one of the substituents R^3 , R^4 , R^5 or R^6 may be transformed to other nitrogen containing substituents under conditions known in the art. For example, alkylation at nitrogen may be performed with an aldehyde under reducing conditions. For acylation the corresponding acyl chloride ($Z = \text{Cl}$) is preferred. Alternatively, an acid anhydride may be used, or acylation may be accomplished with the free acid ($Z = \text{OH}$) under conditions used for amide formation known *per se* in peptide chemistry, e.g. with activating agents for the

carboxy group, such as 1-hydroxybenzotriazole, optionally in the presence of suitable catalysts or co-reagents.

5 Compounds of formula (I) wherein X = NOH may be alkylated allowing access to the corresponding oxime ethers. The reaction conditions leading to this transformation include combinations of weak bases and alkylating agents. Typical bases include metal carbonates or bicarbonates.

10 Reduction of a nitro group in an nitro-substituted aryl or heteroaryl group R or in one of the substituents R³, R⁴, R⁵ or R⁶ to give the corresponding amino group is done, e.g., with iron powder in alcohol or with other reducing agents.

15 A carboxy group in a carboxy-substituted aryl or heteroaryl group R or in one of the substituents R³, R⁴, R⁵ or R⁶ may be amidated under conditions used for amide formation known *per se* in peptide chemistry, e.g. with the corresponding amine and an activating agent for the carboxy group, such as 1-hydroxybenzotriazole, optionally in the presence of suitable catalysts or co-reagents.

20 A bromo or iodo substituent in an aryl or heteroaryl group R or in one of the substituents R³, R⁴, R⁵ or R⁶ may be replaced by phenyl or a phenyl derivative by reaction with a suitable phenylboronic acid in a Suzuki reaction, preferably in a dipolar aprotic solvent such as dimethyl formamide, or in a polar ether, e.g. tetrahydrofuran or dimethoxyethane, in the presence of a soluble palladium(0) or related metal catalyst, for example tetrakis-(triphenylphosphine)palladium.

25 Salts of a compound of formula (I) with a salt-forming group may be prepared in a manner known *per se*. Acid addition salts of compounds of formula (I) may thus be obtained by treatment with an acid or with a suitable anion exchange reagent.

30 Salts can usually be converted to free compounds, e.g. by treating with suitable basic agents, for example with alkali metal carbonates, alkali metal hydrogencarbonates, or alkali metal hydroxides, typically potassium carbonate or sodium hydroxide.

35 It should be emphasized that reactions analogous to the conversions mentioned in this chapter may also take place at the level of appropriate intermediates.

All process steps described here can be carried out under known reaction conditions, preferably under those specifically mentioned, in the absence of or usually in the presence of solvents or diluents, preferably such as are inert to the reagents used and able to dissolve these, in the absence or presence of catalysts, condensing agents or neutralising agents, for example ion exchangers, typically cation exchangers, for example in the H⁺ form, depending on the type of reaction and/or reactants at reduced, normal, or elevated temperature, for example in the range from -100°C to about 190°C, preferably from about -80°C to about 150°C, for example at -80 to +60°C, at -20 to +40°C, at room temperature, or at the boiling point of the solvent used, under atmospheric pressure or in a closed vessel, where appropriate under pressure, and/or in an inert atmosphere, for example under argon or nitrogen.

Salts may be present in all starting compounds and transients, if these contain salt-forming groups. Salts may also be present during the reaction of such compounds, provided the reaction is not thereby disturbed.

At all reaction stages, isomeric mixtures that occur can be separated into their individual isomers, e.g. diastereomers or enantiomers, or into any mixtures of isomers, e.g. racemates or diastereomeric mixtures.

The invention relates also to those forms of the process in which one starts from a compound obtainable at any stage as a transient and carries out the missing steps, or breaks off the process at any stage, or forms a starting material under the reaction conditions, or uses said starting material in the form of a reactive derivative or salt, or produces a compound obtainable by means of the process according to the invention and further processes the said compound *in situ*. In the preferred embodiment, one starts from those starting materials which lead to the compounds described hereinabove as preferred, particularly as especially preferred, primarily preferred, and/or preferred above all.

In the preferred embodiment, a compound of formula (I) is prepared according to or in analogy to the processes and process steps defined in the Examples.

The compounds of formula (I), including their salts, are also obtainable in the form of hydrates, or their crystals can include for example the solvent used for crystallization, i.e. be present as solvates.

New starting materials and/or intermediates, as well as processes for the preparation thereof, are likewise the subject of this invention. In the preferred embodiment, such starting materials are used and reaction conditions so selected as to enable the preferred compounds to be obtained.

5

Starting materials of formula (II) and (III) are known, commercially available, or can be synthesized in analogy to or according to methods that are known in the art.

Pharmaceutical preparations, methods, and uses

10

The present invention relates also to pharmaceutical compositions that comprise a compound of formula (I) as active ingredient and that can be used especially in the treatment of the diseases mentioned at the beginning. Compositions for enteral administration, such as nasal, buccal, rectal or, especially, oral administration, and for

15

parenteral administration, such as intravenous, intramuscular or subcutaneous administration, to warm-blooded animals, especially humans, are especially preferred.

The compositions comprise the active ingredient alone or, preferably, together with a pharmaceutically acceptable carrier. The dosage of the active ingredient depends upon the disease to be treated and upon the species, its age, weight, and individual condition, the individual pharmacokinetic data, and the mode of administration.

20

The present invention relates especially to pharmaceutical compositions that comprise a compound of formula (I), a tautomer, a prodrug or a pharmaceutically acceptable salt, or a hydrate or solvate thereof, and at least one pharmaceutically acceptable carrier.

25

The invention relates also to pharmaceutical compositions for use in a method for the prophylactic or especially therapeutic management of the human or animal body, in particular in a method of treating neoplastic disease, autoimmune disease, transplantation related pathology and/or degenerative disease, especially those mentioned hereinabove.

30

The invention relates also to processes and to the use of compounds of formula (I) thereof for the preparation of pharmaceutical preparations which comprise compounds of formula (I) as active component (active ingredient).

A pharmaceutical composition for the prophylactic or especially therapeutic management of a neoplastic disease, autoimmune disease, transplantation-related pathology and/or degenerative disease, of a warm-blooded animal, especially a human or a commercially useful mammal requiring such treatment, comprising a novel compound of formula (I) as
5 active ingredient in a quantity that is prophylactically or especially therapeutically active against the said diseases, is likewise preferred.

The pharmaceutical compositions comprise from approximately 1% to approximately 95% active ingredient, single-dose administration forms comprising in the preferred
10 embodiment from approximately 20% to approximately 90% active ingredient and forms that are not of single-dose type comprising in the preferred embodiment from approximately 5% to approximately 20% active ingredient. Unit dose forms are, for example, coated and uncoated tablets, ampoules, vials, suppositories, or capsules. Further dosage forms are, for example, ointments, creams, pastes, foams, tinctures,
15 lip-sticks, drops, sprays, dispersions, etc. Examples are capsules containing from about 0.05 g to about 1.0 g active ingredient.

The pharmaceutical compositions of the present invention are prepared in a manner known *per se*, for example by means of conventional mixing, granulating, coating,
20 dissolving or lyophilizing processes.

Preference is given to the use of solutions of the active ingredient, and also suspensions or dispersions, especially isotonic aqueous solutions, dispersions or suspensions which, for example in the case of lyophilized compositions comprising the active ingredient alone
25 or together with a carrier, for example mannitol, can be made up before use. The pharmaceutical compositions may be sterilized and/or may comprise excipients, for example preservatives, stabilizers, wetting agents and/or emulsifiers, solubilizers, salts for regulating osmotic pressure and/or buffers and are prepared in a manner known *per se*, for example by means of conventional dissolving and lyophilizing processes. The said
30 solutions or suspensions may comprise viscosity-increasing agents, typically sodium carboxymethylcellulose, carboxymethylcellulose, dextran, polyvinylpyrrolidone, or gelatins, or also solubilizers, e.g. Tween 80® (polyoxyethylene(20)sorbitan mono-oleate).

Suspensions in oil comprise as the oil component the vegetable, synthetic, or semi-
35 synthetic oils customary for injection purposes. In respect of such, special mention may be

made of liquid fatty acid esters that contain as the acid component a long-chained fatty acid having from 8 to 22, especially from 12 to 22, carbon atoms. The alcohol component of these fatty acid esters has a maximum of 6 carbon atoms and is a monovalent or polyvalent, for example a mono-, di- or trivalent, alcohol, especially glycol and glycerol. As
5 mixtures of fatty acid esters, vegetable oils such as cottonseed oil, almond oil, olive oil, castor oil, sesame oil, soybean oil and groundnut oil are especially useful.

The manufacture of injectable preparations is usually carried out under sterile conditions, as is the filling, for example, into ampoules or vials, and the sealing of the containers.

10 Suitable carriers are especially fillers, such as sugars, for example lactose, saccharose, mannitol or sorbitol, cellulose preparations, and/or calcium phosphates, for example tricalcium phosphate or calcium hydrogen phosphate, and also binders, such as starches, for example corn, wheat, rice or potato starch, methylcellulose, hydroxypropyl
15 methylcellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone, and/or, if desired, disintegrators, such as the above-mentioned starches, also carboxymethyl starch, crosslinked polyvinylpyrrolidone, alginic acid or a salt thereof, such as sodium alginate. Additional excipients are especially flow conditioners and lubricants, for example silicic acid, talc, stearic acid or salts thereof, such as magnesium or calcium stearate,
20 and/or polyethylene glycol, or derivatives thereof.

Tablet cores can be provided with suitable, optionally enteric, coatings through the use of, inter alia, concentrated sugar solutions which may comprise gum arabic, talc, polyvinylpyrrolidone, polyethylene glycol and/or titanium dioxide, or coating solutions in suitable
25 organic solvents or solvent mixtures, or, for the preparation of enteric coatings, solutions of suitable cellulose preparations, such as acetylcellulose phthalate or hydroxypropylmethylcellulose phthalate. Dyes or pigments may be added to the tablets or tablet coatings, for example for identification purposes or to indicate different doses of active ingredient.

30 Pharmaceutical compositions for oral administration also include hard capsules consisting of gelatin, and also soft, sealed capsules consisting of gelatin and a plasticizer, such as glycerol or sorbitol. The hard capsules may contain the active ingredient in the form of granules, for example in admixture with fillers, such as corn starch, binders, and/or
35 glidants, such as talc or magnesium stearate, and optionally stabilizers. In soft capsules,

the active ingredient is preferably dissolved or suspended in suitable liquid excipients, such as fatty oils, paraffin-oil-or-liquid polyethylene glycols or fatty acid esters of ethylene or propylene glycol, to which stabilizers and detergents, for example of the polyoxy-ethylene sorbitan fatty acid ester type, may also be added.

5

Pharmaceutical compositions suitable for rectal administration are, for example, suppositories that consist of a combination of the active ingredient and a suppository base. Suitable suppository bases are, for example, natural or synthetic triglycerides, paraffin hydrocarbons, polyethylene glycols or higher alkanols.

10

For parenteral administration, aqueous solutions of an active ingredient in water-soluble form, for example of a water-soluble salt, or aqueous injection suspensions that contain viscosity-increasing substances, for example sodium carboxymethylcellulose, sorbitol and/or dextran, and, if desired, stabilizers, are especially suitable. The active ingredient, optionally together with excipients, can also be in the form of a lyophilizate and can be made into a solution before parenteral administration by the addition of suitable solvents.

15

Solutions such as are used, for example, for parenteral administration can also be employed as infusion solutions.

20

Preferred preservatives are, for example, antioxidants, such as ascorbic acid, or microbicides, such as sorbic acid or benzoic acid.

25

The present invention relates furthermore to a method for the treatment of a neoplastic disease, autoimmune disease, transplantation related pathology and/or degenerative disease, which comprises administering a compound of formula (I) or a pharmaceutically acceptable salt thereof, wherein the radicals and symbols have the meanings as defined above for formula (I), in a quantity effective against said disease, to a warm-blooded animal requiring such treatment. The compounds of formula (I) can be administered as such or especially in the form of pharmaceutical compositions, prophylactically or therapeutically, preferably in an amount effective against the said diseases, to a warm-blooded animal, for example a human, requiring such treatment. In the case of an individual having a bodyweight of about 70 kg the daily dose administered is from approximately 0.05 g to approximately 5 g, preferably from approximately 0.25 g to approximately 1.5 g, of a compound of the present invention.

30

35

The present invention relates especially also to the use of a compound of formula (I), or a pharmaceutically acceptable salt thereof, especially a compound of formula (I) which is said to be preferred, or a pharmaceutically acceptable salt thereof, as such or in the form of a pharmaceutical formulation with at least one pharmaceutically acceptable carrier for the therapeutic and also prophylactic management of one or more of the diseases mentioned hereinabove, in particular a neoplastic disease, autoimmune disease, transplantation related pathology and/or degenerative disease.

The preferred dose quantity, composition, and preparation of pharmaceutical formulations (medicines) which are to be used in each case are described above.

The following Examples serve to illustrate the invention without limiting the invention in its scope.

15 Examples

Abbreviations: DMSO = dimethyl sulfoxide; THF = tetrahydrofuran,

Example 1: 4-[1-(3-Bromobenzyl)-1H-benzimidazol-2-yl]-furazan-3-ylamine

20 3-Bromobenzylbromide (0.68 g, 2 mmol) is added to an efficiently stirred suspension of 4-(1H-benzimidazol-2-yl)-furazan-3-ylamine (0.5 g, 2 mmol) [A.V. Sergievskii, O.A. Krasnoshek, S.F.Mel'nikova, I.V.Tselinskii, Russian Journal of Organic Chemistry, 2002, 38, 915-917] and potassium carbonate (0.85 g, 6 mmol) in dry N,N-dimethylformamide (10 ml) at 0-5°C. After 30 minutes the reaction mixture is diluted with diethyl ether and the organic phase is washed repeatedly with brine. Drying of the solvent, filtering and evaporation of the solvent under reduced pressure gives the title compound in crude form. The pure compound is obtained after chromatography on silicagel; ¹H-NMR (400MHz, CDCl₃): 7.86 (m, 1H); 7.38 (m, 4H); 7.32 (s, 1H); 7.06 (t, 1H); 7.05 (d, 1H); 5.90 (s, 4H); m.p. 174-176°C.

Example 2: 4-[1-(3-phenylbenzyl)-1H-benzimidazol-2-yl]-furazan-3-ylamine

35 Tetrakis(triphenylphosphine)palladium (7 mg) is added to a degassed solution of 4-[1-(3-bromobenzyl)-1H-benzimidazol-2-yl]-furazan-3-ylamine (0.05 g, 0.13 mmol), phenylboronic acid (0.16 g, 0.13 mmol) in toluene (10 ml) and aqueous sodium carbonate (10 ml of a 2 M solution). The suspension is heated under an atmosphere of argon for

24 hours. The product is extracted using ethyl acetate. The solvent is dried and evaporated, and the residue purified by column-chromatography to yield the title compound as a colorless solid; ¹H-NMR (400MHz, CDCl₃): 7.90 (m, 1H); 7.49-7.32 (m, 10H); 7.10-7.09 (m, 2H); 6.00-5.90 (m, 4H); m.p. 146-148°C.

5

Example 3: 4-[1-(3-Nitrobenzyl)-1H-benzimidazol-2-yl]-furazan-3-ylamine

3-Nitrobenzylbromide (0.1 g, 0.4 mmol) is added to an efficiently stirred suspension of 4-(1H-benzimidazol-2-yl)-furazan-3-ylamine (0.1 g, 0.4 mmol) and potassium carbonate (0.85 g, 6 mmol) in dry N,N-dimethylformamide (10 ml) at room temperature. After 7 hours the reaction mixture is diluted with diethyl ether and the organic phase is washed repeatedly with brine. Drying of the solvent, filtering and evaporation of the solvent under reduced pressure gives the title compound in crude form. The pure compound is obtained after triturating the crude product with hexane; ¹H-NMR (400MHz, d⁶-DMSO): 8.11 (t, 2H); 7.89 (d, 1H); 7.78 (d, 1H); 7.58 (t, 2H); 7.41 (t, 2H); 7.02 (s, 2H); 6.10 (s, 2H); m.p. 238°C.

15

Example 4: 4-[1-(3-Aminobenzyl)-1H-benzimidazol-2-yl]-furazan-3-ylamine

A suspension of 4-[1-(3-nitrobenzyl)-1H-benzimidazol-2-yl]-furazan-3-ylamine (0.1 g, 0.29 mmol) and iron powder (0.16 g, 2.97 mmol) in ethanol (10 ml) is refluxed for 3 hours. The mixture is filtered over a bed of celite and the residue obtained after evaporation of the solvent is purified by chromatography; ¹H-NMR (400MHz, d⁶-DMSO): 7.86 (d, 1H); 7.66 (d, 1H); 7.38 (m, 2H); 7.04 (s, 2H); 6.93 (m, 1H); 6.42-6.24 (m, 3H); 5.82 (s, 2H); 5.20 (s, 2H); m.p. 110°C.

20

Example 5: 4-[1-(3-N-Acetylaminobenzyl)-1H-benzimidazol-2-yl]-furazan-3-ylamine

Acetic anhydride (0.011 ml, 0.1 mmol) is added to a solution of 4-[1-(3-aminobenzyl)-1H-benzimidazol-2-yl]-furazan-3-ylamine (0.03 g, 0.1 mmol) in N,N-dimethylformamide (1 ml) and pyridine (0.01 ml, 0.15 mmol). After stirring the mixture overnight the reaction mixture is partitioned between ethyl acetate and water. The organic layer is separated and dried. Evaporation of the solvent and chromatography on silicagel give the title compound; ¹H-NMR (400MHz, d⁶-DMSO): 9.84 (s, 1H); 7.86 (d, 1H); 7.66 (d, 1H); 7.53 (d, 1H); 7.38 (m, 2H); 7.22 (m, 2H); 7.04 (s, 2H); 6.88 (d, 1H); 5.93 (s, 2H); 1.94 (s, 3H); m.p. 198-202°C.

25

30

35

Example 6: 4-[1-(3-Carboxybenzyl)-1H-benzimidazol-2-yl]-furazan-3-ylamine

3-Bromomethyl-benzoic acid methyl ester (0.11 g, 0.4 mmol) is added to an efficiently stirred suspension of 4-(1H-benzimidazol-2-yl)-furazan-3-ylamine (0.1 g, 0.4 mmol) and potassium carbonate (0.17 g, 1.2 mmol) in dry N,N-dimethylformamide (5 ml) at RT. After heating the mixture for 10 hours at 70°C ethyl acetate is added and the organic phase is washed repeatedly with brine. Drying of the solvent, filtering and evaporation of the solvent under reduced pressure gives the title compound in form of its methyl ester. The free acid is obtained by hydrolyzing the methyl ester with potassium hydroxide.

10 Example 7: 4-[1-(3-(Aminocarbonyl)-benzyl)-1H-benzimidazol-2-yl]-furazan-3-ylamine

To a well stirred solution of 4-[1-(3-carboxybenzyl)-1H-benzimidazol-2-yl]-furazan-3-ylamine (0.064 g, 0.19 mmol) in THF (5 ml) triethylamine (0.019 g, 0.29 mmol) is added. The solution is cooled to 0°C before adding ethyl chloroformate (0.021 g, 0.19 mmol). The reaction mixture is stirred for an additional hour before bubbling ammonia into the reaction solution. The resulting mixture is partitioned between ethyl acetate and water. Separation of the organic phase, drying and purification of the crude product by silica gel chromatography yields the title compound; m.p. 254-256°C.

Example 8: 4-(1-Phenacyl-1H-benzimidazol-2-yl)-furazan-3-ylamine

20 Phenacylbromid (0.1 g, 0.49 mmol) is added to an efficiently stirred suspension of 4-(1H-benzimidazol-2-yl)-furazan-3-ylamine (0.1 g, 0.4 mmol) and potassium carbonate (0.172 g, 1.24 mmol) in dry N,N-dimethylformamide (5 ml) at room temperature. After 4 hours the reaction mixture is diluted with ethyl acetate and the organic phase is washed repeatedly with brine. Drying of the solvent, filtering and evaporation of the solvent under reduced pressure gives the title compound in crude form. The title compound is obtained in pure form by chromatography over silicagel, m.p. 202-204°C.


Example 9: 4-[1-(4-Bromophenacyl)-1H-benzimidazol-2-yl]-furazan-3-ylamine oxime

30 A mixture of 4-[1-(4-bromophenacyl)-1H-benzimidazol-2-yl]-furazan-3-ylamine (0.083 g, 0.21 mmol), sodium bicarbonate (0.021 g, 0.25 mmol) and hydroxylamine hydrochloride (0.014 g, 0.21 mmol) in ethanol (5 ml) is refluxed for 20 hours. Partitioning of the reaction mixture between ethyl acetate and water, separation of the organic phase followed by drying and evaporation of the solvent gives the crude product. Purification by chromatography on silicagel yields the title compound as an E/Z-mixture; m.p. 198-201°C.












Example 10: 4-[1-(2,2-Diethoxy-2-phenylethyl)-1H-benzimidazol-2-yl]-furazan-3-ylamine
4-(1-Phenacyl-1H-benzimidazol-2-yl)-furazan-3-ylamine (0.1 g, 0.3 mmol) is dissolved in HCl / ethanol (2 ml, 8% solution). The mixture is stirred for 30 minutes at room temperature. Addition of diethyl ether and filtration of the resulting precipitate gives the
5 title compound as a colourless solid, m.p. 213-215°C. ¹H-NMR (CDCl₃): 8.22-8.14 (m, 1H); 8.08-8.05 (m, 2H); 7.78-7.50 (m, 6H); 6.25 (s, 2H); 3.72 (q, 4H); 1.24 (t, 6H).

The following compounds were prepared in analogy to Examples 1-10:






















10 Table 1:







15

	R	X	R ⁴ ; R ⁵	Salt	Comment	m.p.
11		bond	H, H	-		
12		bond	H, H	-		172-174°C
13		bond	H, H	-		150°C
14		bond	H, H	-		186-190°C
15		bond	H, H	-		191-194°C
16		bond	H, H	-		140-142°C
17		bond	H, H	-		248°C
18		bond	H, H	-		256-258°C
19		bond	H, F; F, H	-	mixture of isomers	174-176°C
20		bond	H, H	-		214-218°C
21		bond	H, H	-		210-213°C

	R	X	R ⁴ ; R ⁵	Salt	Comment	m.p.
22		bond	H, H	-		oil
23			H, H	-		198-200°C
24		bond	CH ₃ , CH ₃	-		219°C
25		bond	H, COOCH ₃ COOCH ₃ , H	-	mixture of isomers	186°C
26		bond	H, CN CN, H	-	mixture of isomers	210°C
27			H, H	-	E/Z	210°C
28			H, H	-	E/Z	156-158°C
29			H, H	-		160-162°C
30			H, H	-	E/Z	176-178°C
31			H, H	-	E/Z	204-206°C
32			H, H	-	E/Z	206-208°C

	R	X	R ⁴ ; R ⁵	Salt	Comment	m.p.
33			H, H	-		181-184°C
34			H, H	-	E/Z	189-192°C
35			H, H	-	E/Z	160°C
36			H, H	-		226-228°C
37			H, H	-	E/Z	196-198°C
38			H, H	-	E/Z	164-166°C
39			H, H	-		212-214°C
40			H, H	-		198-201°C
41			H, H	-	E/Z	150-154°C
42		— bond —	H, H	-		>250°C
42a		bond	H, H	HCl		> 250°C
43		bond	H, H	-		168-169°C

	R	X	R ⁴ ; R ⁵	Salt		m.p.
44			H, H			168°C
45		bond	H, H			232-234°C
46		bond	H, H			263-264°C

5 General methods for testing of compounds of the invention:

Example 47: Cell cultures and cell lines.

Cell lines are cultured in RPMI-1640 tissue culture medium containing either 5% or 10% fetal calf serum, 0.05 mM 2-mercaptoethanol, 2 mM glutamine and penicillin/streptomycin
 10 50 µg/ml (complete medium) (Sigma, Buchs, Switzerland). General growth conditions are 37°C and 7.5% CO₂.

The following mouse cell lines (either EGFP transfected or not) are being used: A20.2J (ATCC: TIB-208), MC57G (ATCC: CRL-2295).

The following human cell lines (either EGFP transfected or not) are being used: HeLa
 15 (ATCC: CCL-2), KB (ATCC: CCL-17), MCF7 (ATCC: HTB-22), SK-BR-3 (ATCC: HTB-30), SK-Mel 1 (ATCC: HTB-67), SK-Mel 28 (ATCC: HTB-72), PC-3 (ATCC: CRL-1435), SW 480 (ATCC: CCL-228), NCI-H460 (ATCC: HTB-177), NCI-H1792 (ATCC: CRL-5895), HT1080 (ATCC: CCL-21), Jurkat (ATCC: TIB-152), Ramos (ATCC: CRL-1596), Raji (ATCC: CCL-86), H9 (ATCC: HTB-176), Hut78 (ATCC: TIB-161), K562 (ATCC: CCL 243),
 20 HL-60 (ATCC: CCL 240), U-87MG (ATCC: HTB-14), HepG2 (ATCC: HB-8065), U-2 OS (ATCC: HTB-96), Saos-2 (ATCC: HTB-85), U937 (ATCC: CRL 1593), Hs 578T (ATCC: HTB 126), HBL-100 (ATCC: HTB 124), Molt-4 (ATCC: CRL 1582).

Example 48: Primary screening setup

All the manipulations are performed under sterile conditions. The assays are being performed in commercially available 96 or 384 well flat bottom clear microtiter plates (Greiner, Germany) respectively, which are suitable for tissue culture techniques.

- 5 A defined number of EGFP transfected adherent test cells (96 well plates: $10^4 - 10^5$, 384 well plates: $1500 - 2 \cdot 10^4$) are plated out 24 hours before treatment either in 75 μ l (96 well plates) or 60 μ l (384 well plates) complete medium per well in order to ensure appropriate cell spreading. For this purpose a peristaltic pump (e.g. Multidrop by Thermo-Labsystems, Finland) or another suitable device is used. Cells in suspension are plated
10 out according to the same procedure but 1 h prior to treatment. Between seeding out and treatment or addition of compounds the cells are incubated at 37°C under 7.5% CO₂.

Subsequently, the compounds under investigation are added at defined concentrations (40 – 80 μ M in either 25 μ l (96 well plates) or 20 μ l (384 well plates) complete medium

- 15 containing max 4% DMSO) with an appropriate device (e.g. liquid handling system, multi channel pipette etc.) resulting in a final concentration in the test well of 10 – 20 μ M compound in max 1% DMSO.

- Immediately after the addition of the compounds to the cells the zero fluorescence value (t = 0 h) is determined by using a fluorescence microplate reader in order to be able to normalize the fluorescence activities. Afterwards, the test plates are further incubated for
20 a total of 48 h at 37°C under 7.5% CO₂ and are shortly removed only for the purpose of measurement at 8 h, 24 h and 48 h, respectively.

Example 49: Measurement and quantification of the primary screening.

- Relative fluorescence activities of EGFP in compound treated test cells in relation to
25 control cells and cells treated with standard drugs are measured by using a BMG Fluostar microplate fluorescence reader equipped with a filter pair for excitation/emission at 485 nm / 520 nm. The optimum signal to noise ratio is detected by using the time-resolved mode of measurement with a delay of 20 μ s and an integration time over 1 ms. The gain is adjusted in such a way that the control cells produce a fluorescence activity of 90% of
30 the maximum. Kinetics is performed by measuring the relative fluorescence activities at t = 0 h, 8 h, 24 h and 48 h. Crude fluorescence activities are individually normalized for different cell numbers and various optical activities of the test compounds / plate-wells by dividing each value from t = 8 h, 24 h and 48 h by the value of t = 0 h resulting in E(8), E(24) and E(48) values. Subsequently, the E(x) values are further processed by forming
35 the inverse (Q-value) of the products E(8)*E(24)*E(48) which result in numbers > 1 for

apoptotic / necrotic activities of the compounds and numbers < 1 for proliferative activities of the compounds. Controls (untreated) show values similar to 1. Compounds producing Q values > 2 are being considered relevant in terms of apoptotic / necrotic activity and are subsequently tested in the secondary screening setup.

5

Example 50: Secondary screening setup.

All the manipulations are performed under sterile conditions. The assays are being performed in case of adherent cells in commercially available 24 well flat bottom tissue culture plates (Greiner, Germany) and in case of suspension cells in polypropylene tubes (P-tubes) 1.4 ml (Matrix, UK), respectively.

10

Adherent test cells: 2×10^4 – 4×10^4 of EGFP transfected cells in 0.5 ml complete medium are plated out 24 h before treatment. At $t = 0$ the medium is removed and 450 μ l new complete medium is added. Subsequently, 50 μ l complete medium containing the test compound in max. 5% DMSO is added resulting in final concentrations of 20 μ M, 10 μ M, 3 μ M, 1 μ M and 0.3 μ M of the test compounds, respectively. After 48 h incubation the cells are harvested and analyzed with fluorescence activated cell scanning device (FACS Calibur™, BD Biosciences) according to standard procedures.

15

Suspension cells: 10^5 test cells in 450 μ l complete medium are pipetted into P-tubes. 50 μ l complete medium containing the compounds (see adherent cells) is added immediately.

20

After 48 h of incubation the test cells are analyzed directly on a FACSCalibur™.

Example 51: Quantification of the secondary screening.

By monitoring the EGFP fluorescence activity in FL1 on a FACSCalibur™, it is possible to distinguish between proliferating cells, apoptotic cells and necrotic cells within the same cell population. The proliferating cells show a high GFP fluorescence activity, the apoptotic population shows an intermediate fluorescence activity whereas the necrotic cells demonstrate a residual fluorescence activity comparable to mock-transfected cells. Within the CellQuest Software (BD Biosciences) three regions are defined in the histogram: M1 comprising the proliferating cells, M2 comprising the apoptotic cell population and M3 comprising the necrotic cell population. As readout the relative abundance of the cells belonging either to M1, M2 or M3 are expressed. Compounds inducing M2 values $> 50\%$ and M3 values $< 30\%$ are being considered relevant and are further tested and characterized in the tertiary / advanced screening setup.

25

30

Example 52: Tertiary screening setup

A) Hoechst 33342 nuclear staining

This assay is performed in 96 well tissue culture plates. Appropriate number of cells (adherent cells: $3 - 5 \times 10^3$, suspension cells: $8 - 10 \times 10^3$) are being seeded out in 80 μ l complete medium. Adherent cells are incubated for 24 h for proper spreading out before addition of test compounds while suspension cells are immediately treated with test compounds after seeding out. The test compounds are added in 20 μ l complete medium containing max 5% DMSO. The final compound concentrations in the assays are 10 μ M, 3 μ M, 1 μ M and 0.3 μ M, respectively. After 24 h or 48 h incubation at culture conditions, 10 μ l medium containing Hoechst 33342 dye (Sigma B-2261) at 2-5 μ g/ml are added to each well. The assay plates are then further incubated for 30 minutes and subsequently analyzed with a standard inverted fluorescence microscope.

The readout allows the determination of the fraction of apoptotic nuclei as well as other morphological criteria specific for apoptosis as a function of the treatment. Results are indicated in Table 4. The following scores are used: 0 relating to no activity, 1 relating to weak activity comprising less than 50% of the cells and score 2 relating to strong activity comprising more than 50% of the cells.

Table 2: Hoechst 33342 nuclear staining

No	conc	K562		Jurkat		Ramos		HeLa		KB		DM	
		24h	48h	24h	48h	24h	48h	24h	48h	24h	48h	24h	48h
1	10	2	2	2	2	2		2	2	2	2	2	2
	3	2	2			2		2	2	2	2	2	2
	1	2	2			1		2	2	2	2	1	0
	0.3	2	1			0		0	0	0	0	0	0
2	10	2	2	2	2	2		2	2	2		2	2
	3	2	2	1	2	2		1	2	2	2	2	2
	1	1	1	1	1	1		0	0	0	2	0	1
	0.3	0	0	1	0	0		0	0	0	1	0	0
3	10							0	1				
	3							0	0				
	1							0	0				
	0.3							0	0				
4	10							2	2				
	3							0	0				
	1							0	0				
	0.3							0	0				

5	10							2	2				
	3							2	2				
	1							1	1				
	0.3							0	1				
6	10			0	0			0	0				
	3			0	0			0	0				
	1			0	0			0	0				
	0.3			0	0			0	0				
7	10			0	0			0	0				
	3			0	0			0	0				
	1			0	0			0	0				
	0.3			0	0			0	0				
8	10			2	2			2	2				
	3			2	2			2	2				
	1			2	2			2	2				
	0.3			2	2			2	2				
11	10	2	2	1	2			1	2	2	2	1	2
	3	1	1	1	2			1	2	2	2	0	0
	1	1	1	1	2			0	0	0	0	0	0
	0.3	0	0	0	0			0	0	0	0	0	0
12	10	2	2	1	1	2	2	2	2	2	2	2	2
	3	2	2	0	0	2	2	2	2	2	2	2	2
	1	2	2	0	0	1	2	1	2	2	2	2	2
	0.3	1	0	0	0	0	2	0	0	0	0	0	0
13	10	2	2	2	2	2	2	2	2	2	2	2	1
	3	2	2	2	2	2	2	2	2	2	2	2	1
	1	2	2	2	2	2	2	2	2	2	2	2	1
	0.3	1	1	2	2	1	1	2	2	2	2	0	0
14	10			1	2			2	2				
	3			1	1			2	2				
	1			1	0			1	2				
	0.3			0	0			0	0				
15	10	2	2	1	0	2	2	2	2	2	2	2	2
	3	2	2	1	0	2	2	1	2	2	2	2	2
	1	2	1	1	0	1	2	2	0	0	0	0	0
	0.3	0	0	0	0	0	0	0	0	0	0	0	0
16	10	2	2	2	2	2		2	2	2	2	2	2
	3	2	2	2	2	0		2	1	2	2	1	1
	1	1	1	1	1	0		1	0	0	0	0	0
	0.3	0	0	0	0	0		0	0	0	0	0	0
17	10	0	0	0	0			0	1	0	0		
	3	0	0	0	0			0	0	0	0		
	1	0	0	0	0			0	0	0	0		
	0.3	0	0	0	0			0	0	0	0		
18	10	1	1	2	2			1	2			0	1
	3	0	0	2	2			1	1			0	0
	1	0	0	1	2			1	1			0	0
	0.3	0	0	0	1			0	0			0	0

19	10							1	1				
	3							0	0				
	1							0	0				
	0.3							0	0				
21	10			0	0			0	0				
	3			0	0			0	0				
	1			0	0			0	0				
	0.3			0	0			0	0				
22	10							2	2				
	3							0	0				
	1							0	0				
	0.3							0	0				
23	10			2	2			2	2				
	3			2	2			2	2				
	1			2	2			2	2				
	0.3			2	2			2	2				

nd:	Not determined due to self fluorescence of the compound
0:	no effect
1:	weak effect
2:	strong effect

5

10 **B) MTS proliferation assay**

The assay is performed in 96 well tissue culture plates. The cells (range : $1.5 \cdot 10^3 - 10^4$) are seeded out in 80 μ l complete medium 24 h prior to compound treatment. The test compounds are added in 20 μ l complete medium containing max 5% DMSO. The final compound concentrations in the assays are 10 μ M, 3 μ M, 1 μ M and 0.3 μ M, respectively.

- 15 The assay plates are incubated for 72 h at culture conditions. The MTS reagent is prepared according to the manufacturer's protocol (Promega G1111). 20 μ l MTS reagent are added to each well, the assay plates are quickly spun and incubated for another 3 h at culture conditions. Subsequently, the plates are shortly shaken and absorption measured with a microplate-reader at 492 nm. IC₅₀ values are determined by graphical analysis and
- 20 are indicated in the Table 5 in μ M concentration.

Table 3: MTS proliferation assay

No	IC 50				
	Jurkat	K562	HeLa	DM	MRC5
1	1	1	1	2	
2	1	1	2	2	2
3	3		3		3
4	2		2		3
5	2	3	3		3
6	2		2		3
7	3		2		3
8	1		1		2
11		2	2	2	
12	1	1	1	2	2
13	1	1	1	1	1
14	1	1	1	1	2
15	1	2	2	2	3
16	2	2	3	3	3
17	3	3	3		3
18	2	3	3		3
19	2		2		3
20	1		1		2
21	3		3		3
22	3		3		3
23	1		1		1

1:	IC 50 < 1µM
2:	1µM < IC 50 < 10µM
3:	IC 50 > 10µM

5

C) AnnexinV/7-AAD staining

Adherent cells ($1 - 2 \times 10^5$) are 24 h prior to compound treatment seeded into 24 well tissue culture plates. Suspension cells are pipetted into P-tubes immediately before treatment.

- 10 Test compounds are added leading to a final concentrations of 10 µM. After 24 h treatment cells are harvested (in case of adherent cells by trypsinization) and transferred to FACS tubes (BD Biosciences). After centrifugation and removal of the supernatant, 100 µl complete medium containing AnnexinV-GST (10 µg) is added, mixed and incubated at 4°C for 30 minutes. Subsequently, the cells are washed once with medium and incubated
- 15 with 100 µl anti-GST Alexa 488 (Molecular Probes A-11131) in medium diluted 1:500 for 30 minutes at 4°C. Then, cells are washed once and stained with 1 µg/ml 7-aminoactinomycin D (7-AAD) (Molecular Probes A-1310) in 250 µl medium and analyzed on the

FACSCalibur™. AnnexinV is measured in FL1 whereas 7-AAD is measured in FL3. In Table 6 the percentage of positively stained cells after treatment with the respective compounds are indicated. The following scores are used: Score 0 relates to less than 10%, score 1 relates to more than 10% but less than 50% and score 2 relates to more than 50% positivity.

Table 4: AnnexinV/7-AAD staining (10 µM)

No	Jurkat	Ramos	A20	HeLa		
1	1	1	1	1		
2	1	1	1	1		
11	1			1		
12	1	2	1	1		
13	1	2	1	1		
14	1		1	1		
15	1	2	1	1		
16	1	1	1	1		
17	1			1		
18	1			1		

10

0:	< 10%
1:	10% < X < 50%
2:	> 50%

D) PI staining for cell cycle distribution

15 1 - $2 \cdot 10^5$ cells are seeded into 24 well tissue culture plates and incubated for 24 h prior to compound addition. Compounds are added for 24 h in a final concentration of 3 µM or 10 µM. Adherent cells are harvested by trypsinization. The cell suspensions are fixed by adding 2 parts ice cold ethanol 100% while vortexing. Then the samples are stored for > 2 h at -20°C. Subsequently the cells are washed with PBS once and resuspended in

20 250 µl PBS containing 50 µg/ml PI (Calbiochem # 537059), then the samples are incubated at 37°C for 30 minutes and subsequently analyzed on a FACSCalibur™ monitoring linear PI fluorescence activity on FL2. . The readout allows the detection of a possible direct or indirect influence of the tested compounds on the cell cycle. The following events can occur: a) Generation of a subG1 peak indicative for DNA

25 fragmentation, b) increase of the cell population arrested in G2M phase. Both events are scored by 1 for weak and 2 for strong occurrence. 0 indicates no occurrence at all. In Table 7 the influences of several tested compounds are demonstrated.

Table 5: PI staining for cell cycle distribution

No	HeLa 10µM		Jurkat 3µM	
	subG1	G2M	subG1	G2M
1	2	2	1	2
2	1	2	1	2
11	2	1	1	1
12	1	2	1	2
13	1	2	1	2
15	2	2	1	2
16	2	0	0	2

0:	no effect
1:	weak effect
2:	strong effect

5

E) BrdU incorporation (proliferation)

Adherent cells are seeded out at $2 - 4 \times 10^4$ cells/well/ml in 24 well tissue culture plates 24 h prior to treatment. Suspension cells are seeded out at 2×10^5 cells/ml/well in 24 well plates. Compounds are added leading to final concentrations of 3 µM and 10 µM, respectively. Subsequently, BrdU (Molecular Probes #B-23151) at 10 µM final concentration is added and the plates are incubated for 48 h. After the incubation cells are processed according to standard procedures. The detection of the incorporated BrdU is done with the anti-bromodeoxyuridine Mab PRB-1, Alexa Fluor 660 conjugate (Molecular Probes #A-21306). The analysis is performed on a FACSCalibur™ by monitoring the fluorescence activity on FL3. The readout reflects DNA synthesis which is a hallmark for proliferation. The scores are defined the following: 0 for no activity, 1 for a moderate inhibition and 2 for strong inhibition of proliferation. Table 8 indicates inhibitory activities of several tested compounds on proliferation.

20

Table 6: BrdU incorporation

No	HeLa		Jurkat 3µM	
	3µM	10µM	3µM	10µM
1	2		1	

0:	no effect
1:	weak effect
2:	strong effect

5

F) Caspase dependencies

Caspase dependencies are being evaluated by combining the compound treatment with the pan-caspase inhibitor zVAD or its control peptide zFA (ICN Pharmaceuticals # FK009 and FK029, respectively). Both peptides are being used at 20 µM concentration. In case

10 of caspase dependencies a clear inhibition of the specific readout in all apoptosis tests should be detected. By comparing the readout of zVAD and zFA treated samples with the compound control it is possible to detect caspase resp. cystein proteinase dependencies.

In case of inhibition by zVAD but not by zFA a clear caspase dependency is obvious. An inhibition by zVAD as well as by zFA points towards the involvement of cystein

15 proteinases in the apoptotic cascade. Table 9 demonstrates the protease dependency of the nuclear fragmentation visualized by Hoechst 33342 staining.


Example 53: Soft Capsules

5000 soft gelatin capsules, each comprising as active ingredient 0.05 g of one of the
20 compounds of formula (I) mentioned in the preceding Examples, are prepared as follows:
250 g pulverized active ingredient is suspended in 2 liter Lauroglykol® (propylene glycol laurate, Gattefossé S.A., Saint Priest, France) and ground in a wet pulverizer to produce a particle size of about 1 to 3 µm. 0.419 g portions of the mixture are then introduced into soft gelatin capsules using a capsule-filling machine.

25

Claims

1. A compound of formula (I)



5

wherein

- R represents aryl or heteroaryl optionally substituted by up to four substituents independently selected from
- 10 alkyl, cycloalkyl, cycloalkyl-lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, lower alkoxy-lower alkoxy-lower alkyl, halo-lower alkoxy-lower alkyl, acyloxy-lower alkyl, heterocyclyl, heterocyclyl-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl, optionally substituted alkenyl, optionally substituted alkynyl,
- 15 hydroxy, lower alkoxy, optionally substituted alkenyloxy, optionally substituted alkinyloxy, cycloalkoxy, halo-lower alkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, heterocyclyloxy, heterocyclyl-lower alkoxy, optionally substituted phenyloxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy, optionally substituted heteroaryl-lower alkoxy, sulfamoyloxy, carbamoyloxy, lower
- 20 alkylcarbonyloxy, amino, monoalkylamino, dialkylamino, aminocarbonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylcarbonylamino wherein heterocyclyl is bound *via* a nitrogen atom, aminosulfonylamino wherein each of the two amino groups is optionally substituted by
- 25 alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylsulfonylamino wherein heterocyclyl is bound *via* a nitrogen atom, lower alkoxycarbonylamino, lower alkylcarbonylamino wherein alkyl is optionally substituted by one or two substituents selected from optionally substituted phenyl, guanidyl, halogen, cyano, alkoxy, optionally substituted phenoxy,

- alkylmercapto and optionally substituted amino; lower alkenylcarbonylamino wherein alkenyl is optionally substituted by one or two substituents selected from lower alkyl, halo-lower alkyl, optionally substituted phenyl, halogen, cyano, alkoxy and optionally substituted amino; amino-lower alkyl or amino-lower alkylamino, wherein the nitrogen
- 5 atom is unsubstituted or substituted by one or two substituents selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl,
- 10 lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclylcarbonyl, carboxy, lower alkoxycarbonyl, hydroxy-lower alkoxycarbonyl, lower alkoxy-lower alkoxycarbonyl, optionally substituted phenyl-lower alkoxycarbonyl, cyano, lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl; halo-lower
- 15 alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, and nitro; and wherein two adjacent substituents together with the atoms of aryl or heteroaryl may form a 5 or 6 membered carbocyclic or heterocyclic ring;
- 20 X represents a bond; oxygen; a group C=Y, wherein Y stands for oxygen, nitrogen substituted by hydroxy, alkoxy or optionally substituted amino; an ethylenediyl group optionally substituted by alkyl; or a group CR⁷R⁸;
- R¹ and R², independently of each other, represent hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, optionally substituted arylalkyl, optionally substituted heteroarylalkyl, hydroxyalkyl,
- 25 alkoxyalkyl, hydroxyalkoxyalkyl, alkoxyalkoxyalkyl, cyanoalkyl, optionally substituted alkenyl, optionally substituted alkynyl, or lower alkylcarbonyl wherein lower alkyl is optionally substituted by one or two substituents selected from aryl, optionally substituted amino, alkoxy and aryloxy,
- 30 R³, R⁴, R⁵ and R⁶, independently of each other, represent hydrogen, lower alkyl, halo-lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, lower alkoxy-lower alkoxy-lower alkyl, halo-lower alkoxy-lower alkyl, heterocyclyl, heterocyclyl-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower

- alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl, optionally substituted alkenyl, optionally substituted alkynyl, hydroxy, lower alkoxy, halo-lower alkoxy, cycloalkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, heterocyclyloxy, heterocyclyl-lower alkoxy,
5. optionally substituted phenyloxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy, optionally substituted heteroaryl-lower alkoxy, amino, carbamoyl, sulfamoyl, amino-lower alkyl or amino-lower alkylamino, wherein in each case the nitrogen atom is unsubstituted or substituted by one or two substituents selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower
- 10 alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl,
- lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally
- 15 substituted heteroarylcarbonyl, heterocyclylcarbonyl, carboxy, lower alkoxycarbonyl, hydroxy-lower alkoxycarbonyl, lower alkoxy-lower alkoxycarbonyl, optionally substituted phenyl-lower alkoxycarbonyl, cyano, lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl, halo-lower alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower
- 20 alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, or nitro,

or R³ and R⁴, R⁴ and R⁵, or R⁵ and R⁶ together with the atoms of the phenyl ring form a 5 or 6 membered carbocyclic or heterocyclic ring;

- 25 R⁷ represents hydrogen, lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, lower alkenyl, lower alkynyl, optionally substituted phenyl, lower alkoxy, lower alkenyloxy, lower alkynyloxy; R⁸ represents hydrogen, lower alkyl, hydroxy, lower alkoxy or lower alkenyloxy, or R⁷ and R⁸ together with the carbon they are bound to form a 5 or 6 membered carbocyclic or heterocyclic ring;

30

and salts thereof;

for use as medicaments.

2. A compound of formula (I) according claim 1, wherein

R represents aryl or heteroaryl optionally substituted by up to four substituents independently selected from

- 5 alkyl, cycloalkyl, cycloalkyl-lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, lower alkoxy-lower alkoxy-lower alkyl, halo-lower alkoxy-lower alkyl, acyloxy-lower alkyl, heterocyclyl, heterocyclyl-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl, optionally substituted alkenyl, optionally substituted alkynyl,
- 10 hydroxy, lower alkoxy, optionally substituted alkenyloxy, optionally substituted alkynyloxy, cycloalkoxy, halo-lower alkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, heterocyclyoxy, heterocyclyl-lower alkoxy, optionally substituted phenyloxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy, optionally substituted heteroaryl-lower alkoxy, sulfamoyloxy, carbamoyloxy, lower
- 15 alkylcarbonyloxy, amino, monoalkylamino, dialkylamino, aminocarbonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylcarbonylamino wherein heterocyclyl is bound *via* a nitrogen atom, aminosulfonylamino wherein each of the two amino groups is optionally substituted by
- 20 alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylsulfonylamino wherein heterocyclyl is bound *via* a nitrogen atom, lower alkoxycarbonylamino, lower alkylcarbonylamino wherein alkyl is optionally substituted by one or two substituents selected from optionally substituted phenyl, guanidyl, halogen, cyano, alkoxy, optionally substituted phenoxy, alkylmercapto and optionally substituted amino; lower alkenylcarbonylamino wherein
- 25 alkenyl is optionally substituted by one or two substituents selected from lower alkyl, halo-lower alkyl, optionally substituted phenyl, halogen, cyano, alkoxy and optionally substituted amino; amino-lower alkyl or amino-lower alkylamino, wherein the nitrogen atom is unsubstituted or substituted by one or two substituents selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally
- 30 substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl, lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclylcarbonyl,

carboxy, lower alkoxycarbonyl, hydroxy-lower alkoxycarbonyl, lower alkoxy-lower alkoxycarbonyl, optionally substituted phenyl-lower alkoxycarbonyl, cyano, lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl, halo-lower alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, and nitro; and wherein two adjacent substituents together with the atoms of aryl or heteroaryl may form a 5 or 6 membered carbocyclic or heterocyclic ring;

10 X represents a bond or a group C=Y, wherein Y stands for oxygen;

R¹ and R², independently of each other, represent hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, optionally substituted arylalkyl, optionally substituted heteroarylalkyl, hydroxyalkyl, alkoxyalkyl, hydroxyalkoxyalkyl, alkoxyalkoxyalkyl, cyanoalkyl, optionally substituted alkenyl, optionally substituted alkynyl, or lower alkylcarbonyl wherein lower alkyl is optionally substituted by one or two substituents selected from aryl, optionally substituted amino, alkoxy and aryloxy;

20 R³, R⁴, R⁵ and R⁶, independently of each other, represent hydrogen, lower alkyl, halo-lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, lower alkoxy-lower alkoxy-lower alkyl, halo-lower alkoxy-lower alkyl, heterocyclyl, heterocyclyl-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl, optionally substituted alkenyl, optionally substituted alkynyl, hydroxy, lower alkoxy, halo-lower alkoxy, cycloalkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, heterocyclyloxy, heterocyclyl-lower alkoxy, 25 optionally substituted phenyloxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy, optionally substituted heteroaryl-lower alkoxy, amino, carbamoyl, sulfamoyl, amino-lower alkyl or amino-lower alkylamino, wherein in each case the nitrogen atom is unsubstituted or substituted by one or two substituents 30 selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl,

lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclylcarbonyl, carboxy, lower alkoxy carbonyl, hydroxy-lower alkoxy carbonyl, lower alkoxy-lower alkoxy carbonyl, optionally substituted phenyl-lower alkoxy carbonyl, cyano,

5 lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl, halo-lower alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, or nitro,

or R³ and R⁴, R⁴ and R⁵, or R⁵ and R⁶ together with the atoms of the phenyl ring form a 5

10 or 6 membered carbocyclic or heterocyclic ring;

and pharmaceutically acceptable salts thereof; for use as medicaments.

3. A compound of formula (I) according claim 1, wherein

15 R represents phenyl, naphthyl, thienyl, furyl, thiazolyl, oxadiazolyl, thiadiazolyl, imidazolyl, pyrazolyl, pyridinyl, pyrimidinyl, benzothienyl, benzofuryl, indolyl, benzisoxazolyl, each optionally substituted by up to four substituents independently selected from

20 alkyl, cycloalkyl, cycloalkyl-lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, lower alkoxy-lower alkoxy-lower alkyl, halo-lower alkoxy-lower alkyl, acyloxy-lower alkyl, heterocyclyl, heterocyclyl-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl, optionally substituted alkenyl, optionally substituted alkynyl, hydroxy, lower alkoxy, optionally substituted alkenyloxy, optionally substituted alkynyloxy,

25 cycloalkoxy, halo-lower alkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, heterocyclyloxy, heterocyclyl-lower alkoxy, optionally substituted phenyloxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy, optionally substituted heteroaryl-lower alkoxy, sulfamoyloxy, carbamoyloxy, lower alkylcarbonyloxy,

30 amino, monoalkylamino, dialkylamino, aminocarbonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylcarbonylamino wherein heterocyclyl is bound *via* a nitrogen atom, aminosulfonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylsulfonylamino wherein heterocyclyl

35 is bound *via* a nitrogen atom, lower alkoxy carbonylamino, lower alkylcarbonylamino

wherein alkyl is optionally substituted by one or two substituents selected from optionally substituted phenyl, guanidyl, halogen, cyano, alkoxy, optionally substituted phenoxy, alkylmercapto and optionally substituted amino; lower alkenylcarbonylamino wherein alkenyl is optionally substituted by one or two substituents selected from lower alkyl, halo-
5 lower alkyl, optionally substituted phenyl, halogen, cyano, alkoxy and optionally substituted amino; amino-lower alkyl or amino-lower alkylamino, wherein the nitrogen atom is unsubstituted or substituted by one or two substituents selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted
10 heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl, lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclylcarbonyl, lower alkylsulfinyl, halo-lower alkylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl,
15 halogen, and nitro;
and wherein two adjacent substituents together with the atoms of aryl or heteroaryl may form a 5 or 6 membered carbocyclic or heterocyclic ring;

X represents a bond or a group C=Y, wherein Y stands for oxygen;

20 R¹ and R², independently of each other, represent hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, optionally substituted arylalkyl, optionally substituted heteroarylalkyl, hydroxyalkyl, alkoxyalkyl, hydroxyalkoxyalkyl, alkoxyalkoxyalkyl, optionally substituted alkenyl, optionally substituted alkynyl, lower alkylcarbonyl wherein lower alkyl is optionally
25 substituted by one or two substituents selected from aryl, optionally substituted amino or aroyl, or optionally substituted phenylcarbonyl; or R¹ and R² together with the atom they are bound to form heterocyclyl;

R³, R⁴, R⁵ and R⁶, independently of each other, represent hydrogen, lower alkyl, halo-
30 lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, cycloalkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, heterocycliloxy, heterocyclyl-lower alkoxy, optionally substituted phenyloxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy, optionally substituted heteroaryl-lower alkoxy, amino, carbamoyl, sulfamoyl, amino-lower alkyl or amino-lower alkylamino, wherein in
35 each case the nitrogen atom is unsubstituted or substituted by one or two substituents

selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen
5 heterocyclyl,
lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclylcarbonyl,
carboxy, lower alkoxycarbonyl, hydroxy-lower alkoxycarbonyl, lower alkoxy-lower alkoxycarbonyl, optionally substituted phenyl-lower alkoxycarbonyl, cyano,
10 lower alkylsulfinyl, halo-lower alkylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, halogen, or nitro;

or R³ and R⁴, R⁴ and R⁵, or R⁵ and R⁶ together represent methylenedioxy;

15 and pharmaceutically acceptable salts thereof, for use as a medicament.

4. A compound of formula (I) according claim 1, wherein

R represents phenyl, naphthyl, thienyl, furyl, thiazolyl, oxadiazolyl, thiadiazolyl, imidazolyl,
20 pyrazolyl, pyridinyl, pyrimidinyl, benzothienyl, benzofuryl, indolyl, benzisoxazolyl,
optionally substituted by up to four substituents independently selected from
alkyl, halo-lower alkyl, phenyl, optionally substituted heteroaryl, lower alkoxy, optionally substituted alkenyloxy, optionally substituted alkynyloxy, lower alkoxy-lower alkoxy,
amino, monoalkylamino, dialkylamino, aminocarbonylamino wherein each of the two
25 amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl,
heterocyclylcarbonylamino wherein heterocyclyl is bound *via* a nitrogen atom,
aminosulfonylamino wherein each of the two amino groups is optionally substituted by
alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylsulfonylamino wherein heterocyclyl
is bound *via* a nitrogen atom, lower alkoxycarbonylamino, lower alkylcarbonylamino
30 wherein alkyl is optionally substituted by alkoxy or optionally substituted amino; lower alkenylcarbonylamino wherein alkenyl is optionally substituted by alkoxy or optionally substituted amino;
lower alkylsulfinyl, halo-lower alkylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl and halogen;

and wherein two adjacent substituents together with the atoms of aryl or heteroaryl may form a 5 or 6 membered carbocyclic or heterocyclic ring;

X represents a bond or a group C=Y, wherein Y stands for oxygen;

5


R¹ and R², independently of each other, represent hydrogen or lower alkylcarbonyl;

R³, R⁴, R⁵ and R⁶, independently of each other, represent hydrogen, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy, carboxy, lower alkoxycarbonyl, cyano or halogen;

10

and pharmaceutically acceptable salts thereof, for use as a medicament.

5. A compound of formula (I)



15 wherein

R represents aryl or heteroaryl optionally substituted by up to four substituents independently selected from

20 alkyl, cycloalkyl, cycloalkyl-lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, lower alkoxy-lower alkoxy-lower alkyl, halo-lower alkoxy-lower alkyl, acyloxy-lower alkyl, heterocyclyl, heterocyclyl-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl, optionally substituted alkenyl, optionally substituted alkinyl, hydroxy, lower alkoxy, optionally substituted alkenyloxy, optionally substituted alkinyloxy, 25 cycloalkoxy, halo-lower alkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, heterocycliloxy, heterocyclyl-lower alkoxy, optionally substituted phenyloxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy,

optionally substituted heteroaryl-lower alkoxy, sulfamoyloxy, carbamoyloxy, lower alkylcarbonyloxy,

- amino, monoalkylamino, dialkylamino, aminocarbonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl,
- 5 heterocyclylcarbonylamino wherein heterocyclyl is bound *via* a nitrogen atom, aminosulfonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylsulfonylamino wherein heterocyclyl is bound *via* a nitrogen atom, lower alkoxycarbonylamino, lower alkylcarbonylamino wherein alkyl is optionally substituted by one or two substituents selected from optionally
- 10 substituted phenyl, guanidyl, halogen, cyano, alkoxy, optionally substituted phenoxy, alkylmercapto and optionally substituted amino; lower alkenylcarbonylamino wherein alkenyl is optionally substituted by one or two substituents selected from lower alkyl, halo-lower alkyl, optionally substituted phenyl, halogen, cyano, alkoxy and optionally substituted amino; amino-lower alkyl or amino-lower alkylamino, wherein the nitrogen
- 15 atom is unsubstituted or substituted by one or two substituents selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl,
- 20 lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclylcarbonyl, carboxy, lower alkoxycarbonyl, hydroxy-lower alkoxycarbonyl, lower alkoxy-lower alkoxycarbonyl, optionally substituted phenyl-lower alkoxycarbonyl, cyano, lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl, halo-lower
- 25 alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, and nitro; and wherein two adjacent substituents together with the atoms of aryl or heteroaryl may form a 5 or 6 membered carbocyclic or heterocyclic ring;
- 30 X represents oxygen; a group C=Y, wherein Y stands for oxygen, nitrogen substituted by hydroxy, alkoxy or optionally substituted amino; an ethylenediyl group optionally substituted by alkyl; or a group CR⁷R⁸;

- R¹ and R², independently of each other, represent hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, optionally substituted arylalkyl, optionally substituted heteroarylalkyl, hydroxyalkyl,
- 35

alkoxyalkyl, hydroxyalkoxyalkyl, alkoxyalkoxyalkyl, cyanoalkyl, optionally substituted alkenyl, optionally substituted alkynyl, or lower alkylcarbonyl wherein lower alkyl is optionally substituted by one or two substituents selected from aryl, optionally substituted amino, alkoxy and aryloxy,

5

R^3 , R^4 , R^5 and R^6 , independently of each other, represent hydrogen, lower alkyl, halo-lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, lower alkoxy-lower alkoxy-lower alkyl, halo-lower alkoxy-lower alkyl, heterocyclyl, heterocyclyl-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower

10

alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl, optionally substituted alkenyl, optionally substituted alkynyl, hydroxy, lower alkoxy, halo-lower alkoxy, cycloalkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, heterocyclyloxy, heterocyclyl-lower alkoxy, optionally substituted phenyloxy, optionally substituted phenyl-lower alkoxy, optionally

15

substituted heteroaryloxy, optionally substituted heteroaryl-lower alkoxy, amino, carbamoyl, sulfamoyl, amino-lower alkyl or amino-lower alkylamino, wherein in each case the nitrogen atom is unsubstituted or substituted by one or two substituents selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl,

20

lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclylcarbonyl,

25

carboxy, lower alkoxycarbonyl, hydroxy-lower alkoxycarbonyl, lower alkoxy-lower alkoxycarbonyl, optionally substituted phenyl-lower alkoxycarbonyl, cyano, lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfanyl, halo-lower alkylsulfanyl, optionally substituted phenylsulfanyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, or nitro,

30

or R^3 and R^4 , R^4 and R^5 , or R^5 and R^6 together with the atoms of the phenyl ring form a 5 or 6 membered carbocyclic or heterocyclic ring;

R^7 represents hydrogen, lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, lower alkenyl, lower alkynyl, optionally substituted phenyl, lower alkoxy, lower alkenyloxy, lower alkynyloxy;

35

R⁸ represents hydrogen, lower alkyl, hydroxy, lower alkoxy or lower alkenyloxy, or R⁷ and R⁸ together with the carbon they are bound to form a 5 or 6-membered carbocyclic or heterocyclic ring;

5 and salts thereof.

6. A compound of formula (I) according claim 5, wherein

10 R represents phenyl, naphthyl, thienyl, furyl, thiazolyl, oxadiazolyl, thiadiazolyl, imidazolyl, pyrazolyl, pyridinyl, pyrimidinyl, benzothienyl, benzofuryl, indolyl, benzoisoxazolyl, optionally substituted by up to four substituents independently selected from alkyl, cycloalkyl, cycloalkyl-lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, lower alkoxy-lower alkoxy-lower alkyl, halo-lower alkoxy-lower alkyl, acyloxy-lower alkyl, heterocyclyl, heterocyclyl-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl, optionally substituted alkenyl, optionally substituted alkynyl, hydroxy, lower alkoxy, optionally substituted alkenyloxy, optionally substituted alkynyloxy, cycloalkoxy, halo-lower alkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, heterocycliloxy, heterocyclyl-lower alkoxy, optionally substituted phenyloxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy, optionally substituted heteroaryl-lower alkoxy, sulfamoyloxy, carbamoyloxy, lower alkylcarbonyloxy, amino, monoalkylamino, dialkylamino, aminocarbonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylcarbonylamino wherein heterocyclyl is bound *v/a* a nitrogen atom, aminosulfonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylsulfonylamino wherein heterocyclyl is bound *via* a nitrogen atom, lower alkoxycarbonylamino, lower alkylcarbonylamino wherein alkyl is optionally substituted by one or two substituents selected from optionally substituted phenyl, guanidyl, halogen, cyano, alkoxy, optionally substituted phenoxy, alkylmercapto and optionally substituted amino; lower alkenylcarbonylamino wherein alkenyl is optionally substituted by one or two substituents selected from lower alkyl, halo-lower alkyl, optionally substituted phenyl, halogen, cyano, alkoxy and optionally substituted amino; amino-lower alkyl or amino-lower alkylamino, wherein the nitrogen atom is unsubstituted or substituted by one or two substituents selected from lower alkyl,

15
20
25
30
35

- cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl, lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclycarbonyl, carboxy, lower alkoxycarbonyl, hydroxy-lower alkoxycarbonyl, lower alkoxy-lower alkoxycarbonyl, optionally substituted phenyl-lower alkoxycarbonyl, cyano, lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl, halo-lower alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, and nitro; and wherein two adjacent substituents together with the atoms of aryl or heteroaryl may form a 5 or 6 membered carbocyclic or heterocyclic ring;
- 15 X represents oxygen; a group $C=Y$, wherein Y stands for oxygen, nitrogen substituted by hydroxy, alkoxy or optionally substituted amino; an ethylenediyl group optionally substituted by alkyl; or a group CR^7R^8 ;
- 20 R^1 and R^2 , independently of each other, represent hydrogen, lower alkylcarbonyl or optionally substituted phenylcarbonyl;
- 25 R^3 , R^4 , R^5 and R^6 , independently of each other, represent hydrogen, lower alkyl, halo-lower alkyl, alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, amino, carbamoyl, sulfamoyl, amino-lower alkyl or amino-lower alkylamino, wherein in each case the nitrogen atom is unsubstituted or substituted by one or two substituents selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylaminocarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl, 30 carboxy, lower alkoxycarbonyl, hydroxy-lower alkoxycarbonyl, lower alkoxy-lower alkoxycarbonyl, optionally substituted phenyl-lower alkoxycarbonyl, cyano, lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl, halo-lower alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, or nitro,

or R³ and R⁴, R⁴ and R⁵, or R⁵ and R⁶ together represent methylenedioxy;

5 R⁷ represents hydrogen, lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, lower alkenyl, lower alkynyl, optionally substituted phenyl, hydroxy, lower alkoxy, lower alkenyloxy, lower alkynyloxy;

R⁸ represents hydrogen, lower alkyl, lower alkoxy or lower alkenyloxy, or

R⁷ and R⁸ together with the carbon they are bound to form a 5 or 6 membered carbocyclic or heterocyclic ring;

10 and salts thereof.


7. A compound of formula (I) according claim 5, wherein

15 R represents phenyl, naphthyl, thienyl, furyl, thiazolyl, oxadiazolyl, thiadiazolyl, imidazolyl, pyrazolyl, pyridinyl, pyrimidinyl, benzothienyl, benzofuryl, indolyl, benzoisoxazolyl, optionally substituted by up to four substituents independently selected from alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, lower alkoxy-lower alkoxy-lower alkyl, halo-lower alkoxy-lower alkyl, acyloxy-lower alkyl, heterocyclyl, heterocyclyl-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower
20 alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl, optionally substituted alkenyl, optionally substituted alkynyl, hydroxy, lower alkoxy, optionally substituted alkenyloxy, optionally substituted alkynyloxy, cycloalkoxy, halo-lower alkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, heterocyclyloxy, heterocyclyl-lower alkoxy, optionally substituted
25 phenyloxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy, optionally substituted heteroaryl-lower alkoxy, sulfamoyloxy, carbamoyloxy, lower alkylcarbonyloxy, amino, monoalkylamino, dialkylamino, aminocarbonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl,
30 heterocyclylcarbonylamino wherein heterocyclyl is bound *via* a nitrogen atom, aminosulfonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylsulfonylamino wherein heterocyclyl is bound *via* a nitrogen atom, lower alkoxycarbonylamino, lower alkylcarbonylamino wherein alkyl is optionally substituted by one or two substituents selected from optionally
35 substituted phenyl, guanidyl, halogen, cyano, alkoxy, optionally substituted phenoxy,

- alkylmercapto and optionally substituted amino; lower alkenylcarbonylamino wherein alkenyl is optionally substituted by one or two substituents selected from lower alkyl, halo-lower alkyl, optionally substituted phenyl, halogen, cyano, alkoxy and optionally substituted amino; amino-lower alkyl or amino-lower alkylamino, wherein the nitrogen
- 5 atom is unsubstituted or substituted by one or two substituents selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl,
- 10 lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclcarbonyl, carboxy, lower alkoxy carbonyl, hydroxy-lower alkoxy carbonyl, lower alkoxy-lower alkoxy carbonyl, optionally substituted phenyl-lower alkoxy carbonyl, cyano, lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl; halo-lower
- 15 alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, and nitro; and wherein two adjacent substituents together with the atoms of aryl or heteroaryl may form a 5 or 6 membered carbocyclic or heterocyclic ring;
- 20 X represents a group $C=Y$, wherein Y stands for oxygen, nitrogen substituted by hydroxy, alkoxy or optionally substituted amino; or a group CR^7R^8 ;
- R^1 and R^2 , independently of each other, represent hydrogen or lower alkylcarbonyl;
- 25 R^3 , R^4 , R^5 and R^6 , independently of each other, represent hydrogen, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy, carboxy, lower alkoxy carbonyl, cyano, halogen or nitro;
- R^7 represents hydrogen, lower alkyl, hydroxy or lower alkoxy;
- R^8 represents hydrogen, lower alkyl or lower alkoxy; or
- 30 R^7 and R^8 together represent 1,2-ethylenedioxy or 3-alkoxypropylene-1,2-dioxy;

and salts thereof.

8. A compound of formula (I)



5 wherein

R represents heteroaryl optionally substituted by up to four substituents independently selected from

alkyl, cycloalkyl, cycloalkyl-lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-

10 lower alkyl, lower alkoxy-lower alkoxy-lower alkyl, halo-lower alkoxy-lower alkyl, acyloxy-lower alkyl, heterocyclyl, heterocyclyl-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl, optionally substituted alkenyl, optionally substituted alkynyl, hydroxy, lower alkoxy, optionally substituted alkenyloxy, optionally substituted alkynyloxy,

15 cycloalkoxy, halo-lower alkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, heterocyclioxy, heterocyclyl-lower alkoxy, optionally substituted phenyloxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy, optionally substituted heteroaryl-lower alkoxy, sulfamoyloxy, carbamoyloxy, lower alkylcarbonyloxy,

20 amino, monoalkylamino, dialkylamino, aminocarbonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylcarbonylamino wherein heterocyclyl is bound *via* a nitrogen atom, aminosulfonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylsulfonylamino wherein heterocyclyl

25 is bound *via* a nitrogen atom, lower alkoxy carbonylamino, lower alkylcarbonylamino wherein alkyl is optionally substituted by one or two substituents selected from optionally substituted phenyl, guanidyl, halogen, cyano, alkoxy, optionally substituted phenoxy, alkylmercapto and optionally substituted amino; lower alkenylcarbonylamino wherein

alkenyl is optionally substituted by one or two substituents selected from lower alkyl, halo-lower-alkyl, optionally-substituted phenyl, halogen, cyano, alkoxy and optionally substituted amino; amino-lower alkyl or amino-lower alkylamino, wherein the nitrogen atom is unsubstituted or substituted by one or two substituents selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl, lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclylcarbonyl, carboxy, lower alkoxycarbonyl, hydroxy-lower alkoxycarbonyl, lower alkoxy-lower alkoxycarbonyl, optionally substituted phenyl-lower alkoxycarbonyl, cyano, lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl, halo-lower alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, and nitro; and wherein two adjacent substituents together with the atoms of aryl or heteroaryl may form a 5 or 6 membered carbocyclic or heterocyclic ring;

X represents a bond;

R^1 and R^2 , independently of each other, represent hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, optionally substituted arylalkyl, optionally substituted heteroarylalkyl, hydroxyalkyl, alkoxyalkyl, hydroxyalkoxyalkyl, alkoxyalkoxyalkyl, cyanoalkyl, optionally substituted alkenyl, optionally substituted alkynyl, or lower alkylcarbonyl wherein lower alkyl is optionally substituted by one or two substituents selected from aryl, optionally substituted amino, alkoxy and aryloxy,

R^3 , R^4 , R^5 and R^6 , independently of each other, represent hydrogen, lower alkyl, halo-lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, lower alkoxy-lower alkoxy-lower alkyl, halo-lower alkoxy-lower alkyl, heterocyclyl, heterocyclyl-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl, optionally substituted alkenyl, optionally substituted alkynyl, hydroxy, lower alkoxy, halo-lower alkoxy, cycloalkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, heterocycliloxy, heterocyclyl-lower alkoxy,

optionally substituted phenoxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy, optionally substituted heteroaryl-lower alkoxy, amino, carbamoyl, sulfamoyl, amino-lower alkyl or amino-lower alkylamino, wherein in each case the nitrogen atom is unsubstituted or substituted by one or two substituents selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl,

lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclcarbonyl, carboxy, lower alkoxy carbonyl, hydroxy-lower alkoxy carbonyl, lower alkoxy-lower alkoxy carbonyl, optionally substituted phenyl-lower alkoxy carbonyl, cyano, lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl, halo-lower alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, or nitro,

or R³ and R⁴, R⁴ and R⁵, or R⁵ and R⁶ together with the atoms of the phenyl ring form a 5 or 6 membered carbocyclic or heterocyclic ring;

and salts thereof;

with the exception of compounds of formula (I) wherein R represents 4-pyridyl, 3,5-diaminotriazinyl, 3-amino-5-piperidinotriazinyl, 5-methyl-1,3,4-oxadiazol-2-yl, 5-amino-1,3,4-thiadiazol-2-yl and 1H-2-tetrazolyl, X is a bond, R¹ is hydrogen or acetyl, and R², R³, R⁴, R⁵ and R⁶ are all hydrogen.

9. A compound of formula (I) according claim 8, wherein

R represents thienyl, furyl, thiazolyl, oxadiazolyl, thiadiazolyl, imidazolyl, pyrazolyl, pyridinyl, pyrimidinyl, benzothienyl, benzofuryl, indolyl, benzoisoxazolyl, optionally substituted by up to four substituents independently selected from alkyl, cycloalkyl, cycloalkyl-lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, lower alkoxy-lower alkoxy-lower alkyl, halo-lower alkoxy-lower alkyl, acyloxy-lower alkyl, heterocyclyl, heterocyclyl-lower alkyl, optionally substituted phenyl, optionally

substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl, optionally substituted alkenyl, optionally substituted alkynyl, hydroxy, lower alkoxy, optionally substituted alkenyloxy, optionally substituted alkynyloxy, cycloalkoxy, halo-lower alkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, heterocyclyloxy, heterocyclyl-lower alkoxy, optionally substituted phenyloxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy, optionally substituted heteroaryl-lower alkoxy, sulfamoyloxy, carbamoyloxy, lower alkylcarbonyloxy,

5 amino, monoalkylamino, dialkylamino, aminocarbonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylcarbonylamino wherein heterocyclyl is bound *via* a nitrogen atom, aminosulfonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylsulfonylamino wherein heterocyclyl is bound *via* a nitrogen atom, lower alkoxycarbonylamino, lower alkylcarbonylamino

10 wherein alkyl is optionally substituted by one or two substituents selected from optionally substituted phenyl, guanidyl, halogen, cyano, alkoxy, optionally substituted phenoxy, alkylmercapto and optionally substituted amino; lower alkenylcarbonylamino wherein alkenyl is optionally substituted by one or two substituents selected from lower alkyl, halo-lower alkyl, optionally substituted phenyl, halogen, cyano, alkoxy and optionally substituted amino; amino-lower alkyl or amino-lower alkylamino, wherein the nitrogen atom is unsubstituted or substituted by one or two substituents selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylcarbonyl, or

15 wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl, lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclylcarbonyl, carboxy, lower alkoxycarbonyl, hydroxy-lower alkoxycarbonyl, lower alkoxy-lower alkoxycarbonyl, optionally substituted phenyl-lower alkoxycarbonyl, cyano,

20 lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl, halo-lower alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, and nitro; and wherein two adjacent substituents together with the atoms of aryl or heteroaryl may form a 5 or 6 membered carbocyclic or heterocyclic ring;

25

30

X represents a bond;

R¹ and R², independently of each other, represent hydrogen, lower alkylcarbonyl or optionally substituted phenylcarbonyl;

5

R³, R⁴, R⁵ and R⁶, independently of each other, represent hydrogen, lower alkyl, halo-lower alkyl, alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, amino, carbamoyl, sulfamoyl, amino-lower alkyl or amino-lower alkylamino, wherein in each case the nitrogen atom is unsubstituted or substituted by one or two substituents selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylaminocarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl, carboxy, lower alkoxycarbonyl, hydroxy-lower alkoxycarbonyl, lower alkoxy-lower alkoxycarbonyl, optionally substituted phenyl-lower alkoxycarbonyl, cyano, lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl, halo-lower alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, or nitro,

20 or R³ and R⁴, R⁴ and R⁵, or R⁵ and R⁶ together represent methylenedioxy;

and salts thereof.

10. A compound of formula (I) according claim 8, wherein

25

R represents thienyl, furyl, thiazolyl, oxadiazolyl, thiadiazolyl, imidazolyl, pyrazolyl, pyridinyl, pyrimidinyl, benzothienyl, benzofuryl, indolyl, benzoisoxazolyl, optionally substituted by up to four substituents independently selected from alkyl, halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, lower alkoxy-lower alkoxy-lower alkyl, halo-lower alkoxy-lower alkyl, acyloxy-lower alkyl, heterocyclyl, heterocyclyl-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl-lower alkyl, optionally substituted alkenyl, optionally substituted alkynyl, hydroxy, lower alkoxy, optionally substituted alkenyloxy, optionally substituted alkynyloxy, cycloalkoxy, halo-lower alkoxy, cycloalkyl-lower alkoxy, hydroxy-lower alkoxy, lower

35

alkoxy-lower alkoxy, heterocyclyloxy, heterocyclyl-lower alkoxy, optionally substituted phenyloxy, optionally substituted phenyl-lower alkoxy, optionally substituted heteroaryloxy, optionally substituted heteroaryl-lower alkoxy, sulfamoyloxy, carbamoyloxy, lower alkylcarbonyloxy,

- 5 amino, monoalkylamino, dialkylamino, aminocarbonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylcarbonylamino wherein heterocyclyl is bound v/a a nitrogen atom, aminosulfonylamino wherein each of the two amino groups is optionally substituted by alkyl, alkenyl, alkynyl or alkoxy-lower alkyl, heterocyclylsulfonylamino wherein heterocyclyl
- 10 -is bound v/a a nitrogen atom, lower alkoxycarbonylamino, lower alkylcarbonylamino wherein alkyl is optionally substituted by one or two substituents selected from optionally substituted phenyl, guanidyl, halogen, cyano, alkoxy, optionally substituted phenoxy, alkylmercapto and optionally substituted amino; lower alkenylcarbonylamino wherein alkenyl is optionally substituted by one or two substituents selected from lower alkyl, halo-
- 15 lower alkyl, optionally substituted phenyl, halogen, cyano, alkoxy and optionally substituted amino; amino-lower alkyl or amino-lower alkylamino, wherein the nitrogen atom is unsubstituted or substituted by one or two substituents selected from lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, optionally substituted phenyl, optionally substituted phenyl-lower alkyl, optionally substituted
- 20 heteroaryl, optionally substituted heteroaryl-lower alkyl and lower alkylcarbonyl, or wherein the two substituents on nitrogen form together with the nitrogen heterocyclyl, lower alkylcarbonyl, cycloalkylcarbonyl, optionally substituted phenylcarbonyl, optionally substituted heteroarylcarbonyl, heterocyclylcarbonyl, carboxy, lower alkoxycarbonyl, hydroxy-lower alkoxycarbonyl, lower alkoxy-lower
- 25 alkoxycarbonyl, optionally substituted phenyl-lower alkoxycarbonyl, cyano, lower alkylmercapto, optionally substituted phenylmercapto, lower alkylsulfinyl, halo-lower alkylsulfinyl, optionally substituted phenylsulfinyl, lower alkylsulfonyl, halo-lower alkylsulfonyl, optionally substituted phenylsulfonyl, aralkylsulfonyl, halogen, and nitro; and wherein two adjacent substituents together with the atoms of aryl or heteroaryl may
- 30 form a 5 or 6 membered carbocyclic or heterocyclic ring;

X represents a bond;

R¹ and R², independently of each other, represent hydrogen or lower alkylcarbonyl;

R^3 , R^4 , R^5 and R^6 , independently of each other, represent hydrogen, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy, carboxy, lower alkoxy carbonyl, cyano, halogen or nitro;


and salts thereof.

5

11. A method for the preparation of a compound of formula (I) according to claim 1, wherein

A) a compound of formula (II)


10



(II)

wherein R^1 , R^2 , R^3 , R^4 , R^5 and R^6 are defined as for formula (I), or a derivative thereof with functional groups in protected form and/or a salt thereof, is alkylated with a halide of the formula (III)


15



wherein R is as defined for formula (I) and Z is a nucleophilic leaving group;


20 or

B) a compound of formula (IV)



(IV)

25 wherein R^1 and R^2 are defined as for formula (I) and R' is alkyl, or a derivative thereof with functional groups in protected form and/or a salt thereof, is reacted with an ortho-phenylene diamine of formula (V),



wherein R, R³, R⁴, R⁵ and R⁶ are defined as for formula (I), or a derivative thereof wherein functional groups are in protected form and/or a salt thereof;

5

any protecting groups in a protected derivative of a compound of the formula (I) are removed;

and, if so desired, an obtainable compound of formula (I) is converted into another compound of formula (I), a free compound of formula (I) is converted into a salt, an


10 obtainable salt of a compound of formula (I) is converted into the free compound or another salt, and/or a mixture of isomeric compounds of formula (I) is separated into the individual isomers.

12. A pharmaceutical composition comprising a compound of formula (I) according to
15 claim 1 and a pharmaceutically acceptable carrier.

13. Use a compound of formula (I) according to claim 1, a prodrug or a pharmaceutically acceptable salt of such a compound for the preparation of a pharmaceutical composition for the treatment of a neoplastic disease, autoimmune disease, transplantation related
20 pathology and/or degenerative disease.

Abstract

The invention relates to compounds of formula (I)




5

wherein R represents aryl or heteroaryl, X is a bond, a carbonyl group, a derivative of a carbonyl group or an ethylenediyl group, and the substituents R¹ to R⁶ have the meanings given in the specification, for use as medicaments, to novel compounds of formula (I), to methods of synthesis of such compounds, to pharmaceutical compositions containing compounds of formula (I), to the use of a compounds of formula (I) for the preparation of a pharmaceutical composition for the treatment of neoplastic and autoimmune diseases, and to methods of treatment of neoplastic and autoimmune diseases using such compounds of formula (I) or of pharmaceutical compositions containing same.

15

IB_04_1723



This Page is inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLORED OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REPERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images
problems checked, please do not report the
problems to the IFW Image Problem Mailbox**